

# Morphogenesis of rod shaped sacculi

Tanneke den Blaauwen<sup>1\*</sup>, Miguel A. de Pedro<sup>3</sup>, Martine Nguyen-Distèche<sup>2</sup>, and J. A. Ayala<sup>3</sup>

<sup>1</sup>Molecular Cytology, Swammerdam Institute for Life Sciences, University of Amsterdam, Kruislaan 316, 1098 SM Amsterdam, P.O. Box 194062, 1090 GB Amsterdam, The Netherlands.

<sup>2</sup>Centre d'Ingénierie des Protéines, Institut de Chimie B6a, Université de Liège, B-4000 Sart Tilman, Belgium

<sup>3</sup>Centro de Biología Molecular “Severo Ochoa” CSIC-UAM, Campus de Cantoblanco, 28049 Madrid, Spain

\*Corresponding author:

Tanneke den Blaauwen

<sup>1</sup>Molecular Cytology, Swammerdam Institute for Life Sciences, University of Amsterdam, Kruislaan 316, 1098 SM Amsterdam, P.O. Box 194062, 1090 GB

Amsterdam, The Netherlands. Phone: #31 205255196, Fax: #31 205257934

E-mail: [blaauwen@science.uva.nl](mailto:blaauwen@science.uva.nl)

## Keywords

Bacillus subtilis, Caulobacter crescentus, cell division, divisome, Elongase, Escherichia coli, PBP, peptidoglycan

## Abstract 200 words

For growth and division of rod shaped bacteria, the cylindrical part of the sacculus has to be elongated and two new cell poles have to be synthesized. The elongation is performed by a protein complex, the elongase that inserts disaccharidepentapeptides at a limited number of discrete sites while using the cytoskeletal MreB helix as tracking device. Upon initiation of cell division by positioning of the cytoskeletal Z-ring at mid cell, a switch from dispersed to concentrated local peptidoglycan synthesis occurs. From this point on,

1 peptidoglycan synthesis is for a large part redirected from elongating activity to synthesis  
2 of new cell poles by the divisome. The divisome might be envisioned as an extended  
3 elongase because apart from its basic peptidoglycan synthesis activity, specific functions  
4 have to be added. These are, the conversion from cylinder to sphere, the invagination of  
5 the outer membrane, and the addition of hydrolases that allow separation of the daughter  
6 cells. The elongase and the divisome are dynamic hyperstructures that probably share  
7 part of their proteins. Although this multifunctionality and flexibility form a barrier to the  
8 functional elucidation of its individual subunits, it helps the cells to survive a variety of  
9 emergency situations and to proliferate securely.

## 12 **1. Introduction**

14 The shape of bacteria is maintained by the shape of their peptidoglycan layer, as the  
15 isolated peptidoglycan bag or sacculus retains the morphology of the cell it is isolated  
16 from (Fig.1). Comprehension of the mechanism that allow bacteria to differentiate their  
17 shape into cell poles and straight cylinders, banana-shaped cylinders, spirals, and  
18 American footballs to mention a few variations [1], is therefore a prerequisite for the  
19 understanding of bacterial morphology. To unseal these mechanisms, it is not sufficient  
20 to understand all the enzymatic reactions involved in the biosynthetic pathway of the  
21 sacculus (This issues .....).

22 For the synthesis of a three-dimensional structure also the coordinates of synthesis  
23 and hydrolysis or its topography has to be included to answer the question of where the

1 peptidoglycan is synthesized. Topological information has to be added because the  
2 sacculus is embedded in the bacterial envelope that is separated from its metabolic  
3 compartment the cytoplasm. Lastly, the dimension, time, has to be implicated because of  
4 the time dependent nature of the bacterial cell cycle. This review will deal with the  
5 questions where, when and by which proteins the peptidoglycan layer synthesis takes  
6 place in rod shaped bacteria.

7       Due to the differential affinity of a variety of  $\beta$ -lactams for penicillin-binding  
8 proteins (PBPs), one of the earliest observations has been that rod shaped bacteria could  
9 grow temporarily either as spheres or as filaments. As a consequence construction of the  
10 cylindrical part of the cell, which might be termed elongation, and of the cell poles,  
11 termed division, have become separate events in most studies. Therefore, we will discuss  
12 the two processes in this order, although this separation might not completely justified as  
13 will follow from this review. By default, the Gram-negative *Escherichia coli* will be the  
14 model organism for rod shaped bacteria and when necessary illustrative data from the  
15 Gram-positive *Bacillus subtilis* or the gram-negative *Caulobacter crescentus* will be  
16 used.

## 18 **2. Proteins involved in elongation and their interaction**

19  
20 The biosynthetic reaction pathway to peptidoglycan occurs in three stages. The  
21 cytoplasmic stage takes care of the synthesis of the activated nucleotide cytoplasmic  
22 precursors UDP-*N*-acetyl-glucosamine (UDP-GlcNAc) and UDP-*N*-acetylmuramyl-  
23 pentapeptide (UDPMurNAc-pp) via an enzymatic reaction cascade described elsewhere

1 in this issue. This is followed by the membrane linked lipid cycle of reactions that  
2 translocates the lipid-linked  $\beta$ -(1,4) disaccharide peptidoglycan building-unit known as  
3 lipid II across the cytoplasmic membrane (This issue ....). In the periplasmic stage, these  
4 precursors are inserted into the peptidoglycan layer by the PBPs (This issue...). The  
5 cytoplasmic stage is thought to be identical for elongation and division of the bacterium.  
6 It is in the membrane that the first signs of discrimination of the two steps seem to occur.  
7 Proteins that cause the cell to grow as spheres when they are defect, such as MreB, MreC,  
8 MreD, RodA, and PBP 2, are either associated with the membrane or integrated in the  
9 membrane.

10 **RodA** is a 370 amino acids residues integral membrane protein of 40 kDa [2] that  
11 was discovered as a morphological mutant that caused the cells to grow as spheres [3].  
12 Conditional mutants or deletions of homologues of RodA in *B. subtilis*, *Salmonella*  
13 *typhimurium* and even in the ovoid coccus *Streptococcus thermophiles* result in  
14 spherical growth [4-6]. Because RodA in association with PBP2 seems to be essential for  
15 peptidoglycan synthesis in PBP1B-free membrane preparations of *E. coli* [7], it has been  
16 postulated that RodA could be involved in the translocation of the lipidII peptidoglycan  
17 precursors across the cytoplasmic membrane. Although, it has been established that  
18 proteins must be involved in the lipid-II translocation and that translocation seems to be  
19 coupled to the transglycosylation reaction of the PBPs or monofunctional  
20 transglycosylases [8], the identity of the protein(s) responsible for the translocation is still  
21 a mystery.

22 Mecillinam or amdinocillin (F11060, [9]) specifically inhibits the peptidoglycan  
23 transpeptidase **PBP2** [10, 11] causing spherical growth, a substantial increase in the

diameter of the cells and subsequently cell death. The latter can be prevented by overexpression of the cell division proteins FtsQAZ (See section 3) or by increasing the stringent response transcription regulator ppGpp [5, 12-14]. The interpretation is that PBP2 is essential for the maintenance of the diameter of the cells (see also section 5) and in its absence the diameter and therefore the volume of the cells increases uncontrolled. Consequently, the concentration of FtsZ becomes too low to initiate cell division and new polar peptidoglycan synthesis. The other rod shape determining RodA and Mre proteins are also not essential provided that the strains contain mutations that prevent the uncontrolled increase of their diameter. Such strains can be mecillinam resistant [15], overproduce ppGpp [5], overproduce cell division proteins [16, 17] or are osmotically protected [18]. Because PBP2 has transpeptidase activity only, it will have to associate with at least one class A PBPs with glycosyltransferase and transpeptidase activity and perhaps also a monofunctional glycosyltransferase. Based on *in vivo* crosslinking experiments some evidence exists that the majority of the PBPs are part of two different protein complexes [19].

**MreB** (originally EnvB) mutants that grew as spheres were discovered in 1969 by Normark [20] and in combination with **MreC** and **MreD** by Wachi et al., in 1987 [21, 22]. The fusion of GFP to the *B. subtilis* MreB protein in 2001 ([23] and in other species [24, 25], which showed that MreB (and its paralogues in *B. subtilis*, Mbl and MreBH) localizes as helix underneath the cytoplasmic membrane and the subsequent publication of its actin-like crystal structure [26] focused the attention of the scientific community on these proteins. MreB is a soluble cytoplasmic protein of 37-kDa that polymerizes in an ATP dependent fashion [26, 27]. MreC is a 40-kDa bitopic membrane protein with a

1 large periplasmic domain and MreD is an integral membrane protein of 19-kDa. Bacterial  
2 two-hybrid analysis [28] showed that MreC interacts with itself, presumably forming a  
3 dimer [29], and it interacts with MreD and with MreB [17]. MreB interacts with MreC  
4 but not with MreD (Fig. 2). The association of the three proteins is essential for lateral  
5 growth as depletion of each of these proteins is sufficient for spherical growth [17].  
6 Interestingly, MreB is not able to localize in its helical pattern in spherical cells that are  
7 depleted of RodA, MreC, or MreD [17], but it localizes normally in cells that are  
8 spherical due to inhibition of the function of PBP2 by mecillinam [30]. Inhibition of  
9 MreB by its specific inhibitor A22 [31, 32] or inhibition of PBP2 by mecillinam results in  
10 a change from rod to sphere in the same time window [30] with identical changes in the  
11 mucopeptide composition of the sacculus [33]. These changes are a significant shortening  
12 of the glycan chain length, a 60% decrease in pentapeptides, a 20% increase in  
13 crosslinking and an increase in the percentage of lipoprotein-containing mucopeptides  
14 [33] This clearly points to a relationship between MreB and PBP2. PBP2 of *Caulobacter*  
15 *crenscentus* localizes in a band like pattern that is dependent on the presence of MreB and  
16 MreC [25, 34, 35]. MreC itself was also shown to localize in a helical pattern in the  
17 periplasm [34, 36, 37]. Ramoplanin binds to the reducing ends of the nascent glycan  
18 chains [38, 39]. Application of a sub-inhibitory concentration of fluorescent ramoplanin  
19 or vancomycin to growing cells revealed a helical staining pattern along the side-walls of  
20 *B. subtilis* [40] of *E. coli* [33]. Together these data suggest the presence of a protein  
21 complex consisting of MreBCD, RodA and PBP2 that directs lateral peptidoglycan  
22 synthesis. The helical track provided by the MreBCD most likely ensures the disperse  
23 insertion of new peptidoglycan precursors into the existing lateral cell wall. Like actin,

1 MreB monomers treadmill through MreB filaments by preferential polymerization at one  
2 filament end and depolymerization at the other filaments end [41, 42]. It seems likely that  
3 this dynamic movement is passed on to the precursor inserting protein complex (see also  
4 section 4.1).

5 The lipid II peptidoglycan precursor that is synthesized by the membrane  
6 associated glycosyltransferase **MurG** (37.7 kDa) [43-45] is supposed to be the substrate  
7 of this elongase. It is, therefore, not surprising that in *E. coli* MurG could be co-  
8 immunoprecipitated with MreB and that the lateral localization pattern of MurG was lost  
9 in an MreCD deletion background [46]. Similarly, the integral membrane protein **MraY**  
10 (40 kDa) [47-49] that catalyzes the transfer of the phospho-MurNAc-pp motif of UDP-  
11 MurNAc-pp to the lipid carrier undecaprenyl phosphate, to form the so-called lipid I  
12 peptidoglycan precursor to be used by MurG [50, 51], was shown to co-  
13 immunoprecipitate with MreB and MurG [46]. Most likely, MraY and MurG are part of  
14 the protein complex that inserts the peptidoglycan precursors into the peptidoglycan layer  
15 during elongation of the cells, and which therefore could be termed **elongase** (Fig. 2). In  
16 addition, the elongase should contain the lipid II translocase if the RodA protein does not  
17 provide this function. Alternatively, a combination of the integral membrane proteins  
18 might form the translocase (Fig 2B).

### 21 **3. Proteins involved in cell division and their interactions**

1 **MurG** and MraY are both essential proteins with no alternatives available in *E. coli*.  
2 Therefore, both proteins are the most likely candidates to provide lipid I and lipid II  
3 precursors for the synthesis of the new cell poles. In agreement with this notion MurG  
4 was found to localize as well-separated foci in the cylindrical part of the cell and as  
5 concentrated foci at the site of division in *E. coli* [46] and in *C. crescentus* [52]. The  
6 presence of the mid cell localization of MurG was only visible after assemblage of all cell  
7 division proteins at mid cell in *E. coli* [46] but coincided with the assemblage of the Z-  
8 ring in *C. crescentus* [52]. The observation in *E. coli* does not exclude a somewhat less  
9 abundant activity of MurG at the initiation of cell division that coincides with the  
10 assembly of the Z-ring.

### 12 **3.1. Assembly of the Z-ring**

13 The tubulin homologue FtsZ [53] is a GTPase that is conserved in most of bacteria. This  
14 abundant protein (~5000 copies per cell in *E. coli* [54] and den Blaauwen unpublished)  
15 polymerizes in a GTP dependent fashion. It is a highly dynamic structure. Subunits  
16 continuously exchange with a half-life of 8-9 s [55]. The Z-ring is built from overlapping  
17 segments of protofilaments of on averaging about 30 subunits (~120 nm in length) [56].  
18 FtsZ outside the ring localizes in a helix-like pattern and moves rapidly within this  
19 pattern independently from MreB helix or MinD oscillation (see for a review on the min  
20 system [57]). Thus FtsZ not only forms the Z ring but also is part of a highly dynamic,  
21 potentially helical cytoskeleton [58, 59].



1 The formation of the Z-ring, which is the earliest step in cell division, requires the  
2 presence of FtsA or ZipA that tether FtsZ filaments to the membrane. They interact with  
3 a short conserved sequence of the C-terminal end of FtsZ [60-62]. ZipA binds to the  
4 membrane through an essential N-terminal transmembrane segment (TMS) that is linked  
5 to the FtsZ interacting C-terminal globular domain by an extended linker region [63].

6 FtsA is an actin structural homolog [64]. It binds to the membrane through a conserved  
7 C-terminal amphipathic helix that does not appear to be specific as it can substitute for  
8 that of MinD. FtsA is proposed to bind to the membrane before interacting with FtsZ, this  
9 interaction plays an essential role in cell division [65]. Cardiolipin-rich domains are  
10 present in the septal and on the polar membrane regions of *E. coli* [66, 67]. The  
11 amphipathic helix of FtsA has basic residues on its hydrophilic side and might interact  
12 with acidic cardiolipin (CL) in the septal region analogous to the polar assembly of  
13 MinD, which is promoted by CL [68]. Interestingly, overexpression of MurG increased  
14 the amount of CL per cell 7 fold [69]. This could imply that the MurG attracts CL at mid  
15 cell.

16 A conserved region in subdomain 2B of FtsA responsible for interaction with  
17 FtsZ has been identified by isolation of FtsA mutants that are able to bind to the  
18 membrane but fail to interact with FtsZ [62]. FtsA\* contains a modification of R286 into  
19 W in subdomain 2B and can bypass the requirement for ZipA in the assembly of the Z-  
20 ring and its functioning in cell division by stabilizing the FtsZ ring [70, 71]. Subdomain  
21 1C of FtsA is involved in the recruitment of the other components of the divisome [72].

22 Two non related intracellular nonessential proteins (ZapA in *E. coli* [73, 74] and  
23 EzrA in *B. subtilis* [75, 76]) appear to affect the stability of the Z-ring. A proper ratio

1 between FtsZ to FtsA or ZipA is required for assembly and maintenance of the Z-ring.  
2 FtsE/X is an ABC transporter combination of an integral membrane protein FtsX and a  
3 cytosolic membrane associated protein FtsE [77]. The complex interacts with FtsZ [78]  
4 and allows the cell to assemble the Z-ring under conditions of high osmolarity [79].

### 6 **3.2 Maturation of the divisome.**

7 The Z-ring is required for the localization of other cell division proteins, FtsK, FtsQ,  
8 FtsL, FtsB, FtsW, PBP3, (also called FtsI), and FtsN to form the divisome (also called  
9 septosome) at mid cell. All these proteins localise in a sequential and interdependent  
10 manner at the division site (Fig. 3). Proteins FtsK-FtsN are recruited almost  
11 simultaneously at the division site approximately 17 min after the assembly of the Z-ring  
12 [80, 81]. A mutant that underexpresses S-adenosylmethionine synthase forms filaments  
13 without visible septa in which FtsQ, FtsW, PBP3 or FtsN fail to assemble to the Z-ring.  
14 This result suggests that some methylation is required before the complete divisome can  
15 be assembled [82]. MraW also called YabC might be the methylase. Interestingly, the  
16 gene *mraW* with unknown function is located in the *dcw* cell division gene cluster  
17 upstream from *ftsL* [83].

18 **FtsK** is a multifunctional and multidomain protein. The cytoplasmic C-terminal  
19 domain forms the translocation machine, which is involved in chromosome segregation  
20 [84]. The N-terminal domain is formed of 4 transmembrane segments and is required for  
21 cell division [85]. Expression of *ftsA(R286W)* or overexpression of *ftsQ* allows the cell  
22 with a complete deletion of *ftsK* to survive and divide, although many of these *ftsK* null  
23 cells formed multiseptate chains [70]. The cytoplasmic and transmembrane domains of

1 FtsQ, are sufficient to confer viability to *ftsK* null cells. These data suggest that FtsK is  
2 involved in stability of the cell division machine and has a role in the closure of the pole.  
3 Deletion of *dacA* encoding PBP5/DD-carboxypeptidase can suppress the *ftsK44(ts)*  
4 phenotype which is due to a modification of G80 into A in a transmembrane segment.  
5 This modification might affect the conformation of a periplasmic loop of FtsK in contact  
6 with peptidoglycan [86].

7 **FtsQ, FtsL and FtsB** are bitopic membrane proteins with a small N-terminal  
8 intracellular region, a transmembrane segment and a larger periplasmic domain. The  
9 structure of the extracellular region of *Geobacillus stearothermophilus* FtsQ comprises  
10 three domains [87]. The N-terminal  $\alpha$  domain corresponds to the POTRA domain  
11 predicted on the basis of bioinformatics analysis and is proposed to function as a  
12 molecular chaperone [88]. The N-terminal region and  $\alpha$  domain seem to act  
13 cooperatively to bring FtsQ to midcell [89]. The  $\beta$  domain has a unique 3D fold and  
14 might modulate interactions of the flanking domains. The C-terminal  $\gamma$  domain is  
15 sensitive to proteases and unstructured in the absence of division proteins and  
16 corresponds to the region that interacts with FtsL and FtsB in *E. coli* [44]. FtsQ interacts  
17 with FtsL, FtsB, PBP3, FtsW and FtsN as shown by a two-hybrid system and co-  
18 immunoprecipitations [90-92]. Overproduction of FtsQ inhibits division of cells  
19 producing FtsZ, FtsA or PBP3 mutants. These data suggest that FtsQ could play a role in  
20 regulating new pole synthesis [93, 94].

21 FtsL and FtsB seem to interact in a coiled-coil structure through their periplasmic  
22 domain. [95]. They are dependent on each other for proper localization and depend on  
23 FtsQ for their localization at the division site. A fusion of FtsQ to ZapA is recruited to the

1 Z-ring independently of FtsA and directs the recruitment of FtsL, FtsB and PBP3. It can  
2 also back recruit FtsK indicating that both proteins can interact directly [96]. A complex  
3 of FtsL, FtsB and FtsQ can be co-immunoprecipitated in the absence of FtsK, FtsW and  
4 PBP3, which indicates that these proteins probably form a complex before assembling  
5 into the divisome [97].

6 Expression of ZapA-FtsL or ZapA-FtsB allow the recruitment of FtsW and PBP3  
7 in the absence of FtsQ showing that the complex L-B is sufficient to recruit FtsW and  
8 PBP3. Expression of ZapA-FtsW results in efficient localization of PBP3 in cells  
9 depleted of FtsA or FtsQ. Targeting of the complex FtsW-PBP3 restores localization of  
10 the FtsQ-L-B complex in cells depleted of FtsA. Thus the late proteins appear capable of  
11 associating into pre-assembled complexes (FtsQ-FtsB-FtsL and FtsW-PBP3) within the  
12 cell [98]. As the average number of FtsQ proteins in the cell is approximatively 20-40  
13 and that of PBP3 ~100, limited number of protein sub-complexes can be expected along  
14 the Z-ring.

15 **FtsW** has 10 transmembrane segments (TMSs) and shares 30% identity with  
16 RodA and SpoVE, which are involved in elongation and in *Bacillus* sporulation  
17 respectively. These proteins appear to work in coordination with one class B PBP to  
18 catalyze peptidoglycan polymerization during the cell cycle (see also section 2). The  
19 periplasmic loop of FtsW from residue P368 to P375 plays an important role in the septal  
20 recruitment of PBP3, the E240-A249 periplasmic amphiphilic sequence appears to be a  
21 key element of the functioning of FtsW in the septal peptidoglycan assembly machineries  
22 [59]. The first 75 amino acid residues of FtsW are sufficient to interact with FtsQ [90].

1       The multimodular class B **PBP3** that is specifically involved in septal  
2 peptidoglycan synthesis consists of a short intracellular M1-R23 peptide fused to a F24-  
3 L39 membrane anchor that is linked via a G40-S70 peptide to the non-catalytic module  
4 itself linked to the penicillin-binding module with transpeptidase activity [99, 100]. The  
5 first 56 amino acid residues of PBP3 possess the structural determinants required to target  
6 the protein to the cell division site, and none of the putative protein interacting peptides  
7 present in the N-terminal non-catalytic module are essential for the positioning of the  
8 protein at the division site [94]. By using an *E. coli* two-hybrid system, the first 56 amino  
9 acid residues of PBP3 were shown to interact with FtsW (unpublished data M. Distèche).  
10 Residues 1 to 70 of PBP3 are sufficient to interact with FtsQ, whereas the membrane  
11 bound non-catalytic module appears to interact with FtsL [92].

12       **PBP1B** localizes at the division site and in the cylindrical part of the cell and  
13 interacts directly with PBP3 as shown by using affinity chromatography, surface plasma  
14 resonance, two-hybrid system and immunoprecipitation [101]. Its localization depends on  
15 the physical presence of PBP3 but not on its activity. A  
16 monofunctionalglycosyltransferase (MGT) is able to interact with three constituents of  
17 the divisome, PBP3, FtsW and FtsN in a bacterial two-hybrid assay (unpublished M.  
18 Terrak and M. Distèche). Finally, PBP1C interacts with PBP3 and PBP1B [102] and the  
19 structural protein MipA interacts as well with PBP1B as with the hydrolase MltA [103]  
20 (see this issue x). The localization of these proteins is not yet known.

21       **FtsN** is a bitopic protein with an N-terminal intracellular domain, a TMS and a  
22 periplasmic domain. This region is formed of three short helices (residues 62-123), a long  
23 glutamine-rich, unstructured region (residues 124-242) and a C-terminal domain which

1 binds peptidoglycan but is not essential for cell division [104]. The N-terminal region  
2 (residues 1-62) of FtsN although normally not essential for the protein function can  
3 compensate for a complete lack of FtsK [105].

4 FtsN interacts with PBP3 and PBP1B and it stimulates PBP1B activity, indicating  
5 that it might have a role in the control of the peptidoglycan synthesis of the third stage of  
6 pole synthesis (unpublished Vollmer and Nguyen-Distèche and see section 4.1). A fusion  
7 of the subdomain 1C of FtsA to the *Bacillus subtilis* DivIVA protein is targeted to the cell  
8 pole and is able to recruit FtsN and PBP3 to the poles independently from FtsZ [106].  
9 FtsN is not recruited by prematurely targeted protein complexes FtsQLB or FtsW-PBP3  
10 and required the presence of FtsA at the division site [98]. FtsA mutants that increase the  
11 integrity of the Z-ring can compensate for the loss of FtsN and can partially compensate  
12 for a deletion of FtsK or ZipA [107]. FtsN could play a role in regulating conformational  
13 changes within the divisome subassemblies, which in turn regulate the activity of the Z-  
14 ring. FtsA appears to be conformationally flexible and could be a key modulator of the  
15 divisome function at all stages [107].

16 **AmiC** is a periplasmic amidase that cleaves the septal peptidoglycan and  
17 promotes cell separation. It depends on the presence of FtsN for its localization and it  
18 contains a N-terminal domain that is necessary and sufficient to target the protein at the  
19 division site [108, 109] (see section 4.1). **EnvC**, a periplasmic metallo-endopeptidase  
20 also participates to the splitting of peptidoglycan septum and is located at the division site  
21 [110].

#### 22 23 **4. Rate and topology of peptidoglycan synthesis during the cell cycle**

1

2        Generation and maintenance of proper morphology in rod shaped bacteria  
3 requires a mechanism able to produce a regular cylindrical surface, the lateral  
4 peptidoglycan, that grows in length (elongation) whilst avoiding generation of bending  
5 and torsional forces. Otherwise growth would result in cells of irregular diameter, with  
6 curved or twisting morphologies, shapes often associated with mutational impairment of  
7 some morphogenes [1, 111].

8        At periodical intervals, concomitantly with the onset of cell division, the mode of  
9 PG synthesis changes from a constant diameter "elongative" mode to a decreasing  
10 diameter "constrictive" mode to form the transversal wall at the cell centre, which will in  
11 turn generate the new cell poles as physical separation of the daughter cells proceeds [11,  
12 112-115]. Both elongation and constriction ultimately rely on the local activity of a set of  
13 complexes where precursors, originated in the cytoplasm, become polymerized and  
14 covalently incorporated into the meshwork structure of the sacculus causing its  
15 expansion. However the mechanisms and proteins involved are different enough as to  
16 treat both independently in this discussion.

17

18

#### 19    **4.1 Topology of insertion sites in the lateral wall of sacculi**

20    Whether incorporation of new precursors into the sacculus happened at a defined location  
21 or in a diffuse way was soon decided in favour of the later alternative. Studies performed  
22 in the 60's already were supportive of diffuse incorporation in a relatively large number  
23 of sites [116]. Application of high resolution autoradiography lead to some initial

1 confusion, as some early results were interpreted in favour of zonal growth [117].  
2 However, further technical and experimental improvements confirmed the disperse nature  
3 of precursor insertion into the lateral wall by a series of complementary approaches as: i)  
4 high resolution autoradiography following incorporation of radioactive precursors [118-  
5 122]; ii) evolution of the acceptor-donor radioactivity ratio a parameter that indicates  
6 whether radioactively labeled incoming precursors cross-link preferentially with  
7 themselves or with preexisting material [123, 124]; iii) distribution of peptidoglycan  
8 hydrolytic [123, 125], biosynthetic [126, 127] and morphogenetic [17, 128] activities; iv)  
9 analysis of the peptidoglycan segregation pattern by D-amino acid labelling and  
10 immunomicroscopy [33, 129]; and v) Ramoplanin affinity labeling of precursors [40] and  
11 newly incorporated peptidoglycan [33]. Therefore, that elongation of sacculi occurs by  
12 incorporation of precursors in a number of homogeneously distributed sites covering the  
13 cylindrical surface of the sacculus seems to be a well established fact, at least for the  
14 model rod-like bacteria *E. coli*, *B. subtilis*, and related species. The number of insertion  
15 sites active at any particular time point is likely to be rather small. Although the earlier  
16 estimations suggested "200 or more" [116], later estimates reduced figures to 80-100 sites  
17 per cell based on the kinetics of insertion [130]. Nonetheless, this figure is likely a  
18 considerable over-estimation, in view of the scarcity of some critical components of the  
19 elongation machinery as PBP2 of *E. coli* (20-40 molecules per cell) [11, 127, 131]. The  
20 real nature of the material in the course of insertion is still under debate, in particular  
21 whether new glycan chains are inserted individually [132], as pairs [130] or as triplets  
22 [113]. Recent evidence supports an intrinsic ability of peptidoglycan synthases to produce  
23 pairs of cross-linked glycan strands[133, 134], therefore supporting the idea that glycan



1 strands are likely to be inserted into the sacculus as cross-linked pairs, or higher order  
2 polymers.

3 The recent demonstration that proper bacterial morphogenesis depends on the  
4 dynamics of cytoskeletal proteins as MreB spirals and FtsZ rings [135] supports the  
5 concept of insertion sites as wandering structures that travel around the sacculus whilst  
6 promoting synthesis and insertion of peptidoglycan strands [17, 33, 111, 127, 128].  
7 Actually, the proposal of insertion sites as dynamic entities is anything but new. It was  
8 first proposed more than twenty years ago [130], and was also an intrinsic assumption in  
9 the "three for one" model for growth of the murein sacculus [113].

10 That insertion of new material may happen at any place of the lateral wall with  
11 equal probability doesn't imply that new and old material is fully intermixed. As a matter  
12 of fact, the (sparse) evidence available rather supports generation of mosaic structures  
13 made up of interconnected microdomains of all-new and all-old peptidoglycan [33, 136,  
14 137].

15 In conclusion, our present view is that insertion of precursors into the lateral wall  
16 of the sacculus happens at a limited number of discrete sites, which are highly mobile and  
17 distributed over the surface of the cell in an MreB associated pattern.

18

19

## 20 **4.2 Topology of precursor insertion sites in septal-polar peptidoglycan**

21

22 Cell poles are the product of cell division, and by extension polar peptidoglycan is the  
23 product of septation. Division of the sacculus requires a change in the mode of synthesis

1 of macromolecular peptidoglycan. At periodic intervals insertion of precursors instead of  
2 promoting elongation starts to generate either an invagination of progressively smaller  
3 diameter, or an inwards growing thick transversal septum in Gram negative and Gram  
4 positive bacilli, respectively [11, 113-115, 138]. This change has been associated with  
5 activation of zonal, that is highly localized, insertion since the earliest studies on cell wall  
6 growth; first as a rationalization of the lytic phenotypes observed in response to  
7 antibiotics as penicillin [125, 139], and second by high resolution autoradiography [117,  
8 120-122, 140], and peptidoglycan segregation studies [129, 141]. The later study  
9 exploited the ability of a, still uncharacterized, periplasmic LD-transpeptidase activity  
10 [142] to mediate the exchange of an externally added D-amino acid (D-cysteine) for the  
11 D-ala normally present at the C-terminal position of mucopeptides. After isolation of the  
12 sacculi, the D-Cys are biotinylated and labeled with biotin recognizing gold-labeled  
13 secondary antibodies. When growth of the cells in the presence of D-Cys was followed  
14 by a chase for one or two mass doublings in the absence of D-Cys, the number of gold  
15 particles on the cell poles was still the same compared to unchased samples, whereas the  
16 number of particles in the cylindrical part of the cells was diluted by growth (Fig. 4).  
17 From these results it was confirmed that the cell poles are constructed from new  
18 peptidoglycan and that they become inert after their synthesis [122, 129].

19 Polar peptidoglycan has a very long life, and seems to remain unchanged for  
20 several generations. Neither synthesis nor degradation of peptidoglycan take place in the  
21 poles, which can therefore be considered as stable structures [67]. No chemical or  
22 structural features clearly accountable for the high stability of polar peptidoglycan have  
23 been detected in sacculi. The relatively minor modifications detected are in general

1 smaller than variations due to growth rate or state of growth [143-147]. An indication that  
2 some modification occurs after synthesis of the new cell poles is that the old cell poles do  
3 contain recycled peptidoglycan muropeptides, whereas the newly synthesized poles are  
4 devoid of these peptides (unpublished data Oldrich, den Blaauwen et al.,). Deficiencies in  
5 PBP5, the main DD-carboxypeptidase activity of *E. coli*, have a dramatic influence on the  
6 topology of inert peptidoglycan, in particular when combined with mutations on other  
7 low molecular weight PBPs as PBP7 [148, 149]. These strains often show morphological  
8 alterations as branches, buds and kinks which are always and every time associated to  
9 regions of inert peptidoglycan [150]. How these ectopic regions are generated is under  
10 study, but evidence indicates interference with normal septation as aberrant poles are a  
11 common feature of such strains [150] (unpublished data M.A. de Pedro). An alternative  
12 possibility is that dynamics of cytoskeletal proteins preclude biosynthetic complexes  
13 from polar regions, as suggested by the influence of FtsZ on peptidoglycan synthesis [52,  
14 111], and in particular on the extension of polar inert peptidoglycan regions [33].  
15 Nevertheless, experimental evidence is insufficient to actually settle the issue.

16       The detailed architecture and distribution of insertion sites involved in septal  
17 peptidoglycan synthesis are still poorly known. Activation of septal peptidoglycan  
18 synthesis seems to be a multistep process both regarding assembly of division complexes  
19 [80] and insertion of precursors into the central region of the sacculus [121]. Septal  
20 synthesis can be divided into at least two stages related to the involvement of cell  
21 division proteins; an early stage which requires FtsZ as the only division specific protein,  
22 and a later stage which requires the full complement of division proteins [129].  
23 Activation of zonal insertion at the potential division site requires assembly of the FtsZ

1 ring at the cell centre, but can proceed for a significant period of time in the absence of  
2 later division proteins as FtsQ and FtsI, leading to the synthesis of an annulus of all-new  
3 murein at the potential division site (Fig. 4).

4       Once a particular, still unknown, check-point is reached in FtsQ and FtsI  
5 filaments, septal synthesis apparently stops and the peptidoglycan biosynthetic potential  
6 becomes redirected towards lateral wall growth leaving behind an annulus of inert  
7 peptidoglycan [129]. This early step is likely to represent the PIPS (penicillin insensitive  
8 peptidoglycan synthesis) stage as proposed by Nanninga [121, 151] on the basis of  
9 observations indicative of a penicillin insensitive nature for the initial phase of cell  
10 division. When division is allowed to proceed, septal synthesis goes on promoting the  
11 progressive constriction of the sacculus. Incorporation of precursors on this phase seems  
12 to take place exclusively at the leading edge of the inwards growing constriction, in a  
13 PBP3 dependent manner [121]. The nature of the leading edge is still unknown, however  
14 evidence from *E. coli* mutants defective in N-acetyl-muramyl-L-alanine amidases  
15 supports the idea of the leading edge as a differentiated domain in septal peptidoglycan.  
16 N-acetyl-muramyl-L-alanine amidases are required at the terminal stages of septation  
17 [145, 152]. Defective *amiABC* mutants, form chains of cells separated by incomplete  
18 septa [109, 152, 153]. Electron microscopy analysis of sacculi from such strains revealed  
19 the existence of an electron dense ring structure (Septal Peptidoglycan or SP-ring)  
20 tagging the inwards edge of partially constricted division sites [109, 153, 154] (Fig. 5).  
21 Furthermore SP-rings are associated to areas of inert peptidoglycan, suggesting that polar  
22 peptidoglycan becomes metabolically inert since the very moment of insertion [154].  
23 Apparently SP-rings are able to progress down to a minimum diameter in *amiABC*

mutants, and then stall. Cytokinesis, however, proceeds to the end and cells become independent entities separated by the respective cytoplasmic membranes [152, 153]. Therefore a potential third (late) stage in septation could be marked by the specific requirement for N-acetyl-muramyl-L-alanine amidases activities to allow the closure of the inwards growing septum and terminate synthesis of polar peptidoglycan.

#### **4.3. Rate of PG synthesis**

The rate of peptidoglycan synthesis has been investigated by pulsed incorporation of radioactive peptidoglycan precursors in synchronous growing cells and asynchronous growing cells [121, 155] (see also 4.1). In both cultures the rate of peptidoglycan synthesis was more or less exponential. However when the incorporation of radioactive *meso*-diaminopimelic acid (A<sub>2</sub>pm, see this issue ...) was visualized by autoradiography and electron microscopy, an increase in the rate of peptidoglycan synthesis at the constriction site at the expense of a reduction of 40% in the rate of peptidoglycan synthesis in the cylindrical part of the cell was observed [119, 121]. The incorporated radioactive A<sub>2</sub>pm was visible after autoradiography as black grains on the cell. In the cylindrical part of the cells the grains were dispersedly distributed whereas they were clustered at the site of division. Even in cells that showed no visible constriction, but that were about to initiate division an increase in the number of grains at mid cell was observed. This increase in mid cell peptidoglycan synthesis is dependent on the presence of FtsZ [119] and coincides with the moment at which the Z-ring is assembled [156]. Remarkably, the increase in mid cell peptidoglycan synthesis is not dependent on the

1 presence or the function of PBP3 [121] or probably any of the cell division proteins that  
2 are recruited to the Z-ring in the second step of divisome maturation [33, 80]. As a  
3 consequence, these proteins cannot cause the increased rate of peptidoglycan synthesis at  
4 mid cell and the accompanying reduction in lateral peptidoglycan synthesis rate. Similar  
5 observations have been made with respect to the two modes of peptidoglycan synthesis  
6 (Fig 4). The polar mode of peptidoglycan synthesis is dependent on FtsZ, but not on  
7 PBP3 or FtsQ [129]. Therefore, it seems acceptable to assume that the polar mode of  
8 peptidoglycan synthesis and the increase in the rate of peptidoglycan synthesis coincide  
9 in time and space.

10       The rate of A<sub>2</sub>pm incorporation can be almost completely inhibited either by  
11 blocking PBP1A/B by their specific inhibitor cefsulodin or by blocking PBP2 and PBP3  
12 simultaneously by mecillinam and cephalexin [157]. This suggests that the PBP1s and the  
13 transpeptidases act together in the synthesis of the cell envelope (see also 4.3). Inhibition  
14 of PBP2 in synchronized cells reduced the rate of A<sub>2</sub>pm incorporation for 60% during the  
15 entire cell cycle whereas inhibition of PBP3 by cephalexin reduced the rate of  
16 incorporation especially during the constriction period for 35% [157]. This shows that  
17 lateral peptidoglycan synthesis continues during the constriction period, albeit at a lower  
18 rate as the number of A<sub>2</sub>pm grains in the lateral wall was reduced by 40%. Because the  
19 PBP2 activity was more or less equally inhibited in non-constricting cells as in  
20 constricting cells (Fig. 6), it can be concluded that PBP2 assist in the synthesis of the new  
21 cell poles (see also section 5) whereas PBP3 seems to be restricted to new cell pole  
22 synthesis.

1       The average number of autoradiography  $A_{2pm}$  grains at the division site did not  
2 diminish in cells that had progressed in polar cap synthesis compared to division  
3 initiating cells [121]. These observations suggest that peptidoglycan synthesis at the  
4 leading edge of the constriction site occurs by more or less the same set of peptidoglycan  
5 synthesizing protein complexes as those that were recruited during the initiation of cell  
6 division. These proteins complexes have to become more clustered during the  
7 constriction process when the FtsZ-ring diminishes in diameter. As a consequence of the  
8 constant rate of polar peptidoglycan synthesis, the Z-ring diameter decreases with a rate  
9 to the power of two. This is in agreement with the relative small fraction of deeply  
10 constricting cells in a population.

#### 11 12 **4.4. Are specific proteins responsible for the local insertion and increase in PG** 13 **synthesis at the initiation of cell division?**

14       The local increase in peptidoglycan synthesis is dependent on the presence of  
15 FtsZ as was discussed above, but not on the proteins that assemble onto the ring in a later  
16 stage [80]. It is accompanied by a lengthening of the glycan stands [144] and the  
17 synthesis of a band of inert peptidoglycan at the future site of division [129]. Curiously  
18 this initiation of constriction, which coincided with Z-ring assembly, could not be  
19 inhibited by penicillins such as cefsoludin, mecillinam, furazlocilin, and cephalexin and  
20 moenomycin [157]. The presence of a penicillin insensitive peptidoglycan synthetase  
21 (PIPS) that interacts with FtsZ was postulated [140] to be responsible for the initiation of  
22 polar peptidoglycan synthesis. Could any of the other known peptidoglycan synthesizing  
23 proteins be part of PIPS? The third class A PBP of *Escherichia coli*, PBP1C, is

1 insensitive to the majority of the penicillins and after immobilization on sepharose it was  
2 able to retain PBP1B, PBP3, and MltA [102]. It could also retain PBP2 provided that the  
3 periplasmic fraction was added to the solubilised membranes [102, 103, 158]. This  
4 suggests that PBP1C is at least part of the complex responsible for polar peptidoglycan  
5 synthesis and that it could represent PIPS. Because PBP1C is reported to bind to  
6 moenomycin coupled agarose it cannot completely represent PIPs. The monofunctional  
7 glycosyl transferase (Mgt) of *E. coli* is insensitive to moenomycin [159] and interacts in a  
8 bacterial two hybrid system with PBP3, FtsW and FtsN (unpublished data Distèche,  
9 Terrak et al.,). The combination of PBP1C and Mgt could provide all the enzymatic  
10 activities for PIPS. However, deletion of either or both proteins simultaneously has no  
11 effect on the morphology of the cells [102]. The penicillin insensitive LD-transpeptidase  
12 of *E. coli* [142] seems an unlikely candidate as it was shown to be involved in the  
13 attachment of the Braun lipoproteine [160]. Since PBP2 is involved in cell pole synthesis  
14 [127], it is possible that PBP2 is able to initiate cell division when either PBP3 or  
15 PBP1A/B are inhibited. If this would be the case, the peptidoglycan synthetases do not  
16 provide any specificity to explain the switch from lateral to polar peptidoglycan  
17 synthesis. Specificity would have to be found in, for instance, the activity of the  
18 hydrolases such as the carboxypeptidases or in the compartmentalization that the  
19 cytoskeleton of the FtsZ-ring and the MreB helix somehow seems to provide. The  
20 occurrence of a switch in the mode of peptidoglycan synthesis that leads to new cell pole  
21 synthesis is firmly established, but its nature, as is outlined above, is very poorly  
22 understood. Future studies might need to include not only the proteins involved in the  
23 synthesis of the peptidoglycan layer, but also the physical arrangement of the



1 biosynthetic complexes as function of the cell cycle and the metabolic and signaling state  
2 of the cells.

### 3 4 5 **5. Is the cellular localization of PBPs related to their function?**

6  
7 The PBPs (see also this issue ...) are responsible for the periplasmic stage of  
8 peptidoglycan synthesis by inserting the lipid II peptidoglycan precursors in the existing  
9 peptidoglycan layer during length growth or by synthesizing new peptidoglycan during  
10 cell division (section 4). Although some of the PBPs have a unique function, many are  
11 redundantly present in most bacterial species investigated thus far. Their specific  
12 function, apart from their enzymatic activity, is not very well known. The exceptions are  
13 the class B high molecular weight PBPs that have transpeptidase activity. In *E. coli* these  
14 are PBP2 and PBP3, which could easily be characterized as essential for length growth  
15 and division [11, 161, 162] because their inhibition has such a dramatic effect on cell  
16 morphology (i.e. conversion to spheres or filaments, respectively). Deletion of the genes  
17 encoding for PBPs has not provided clear-cut information on their function, as PBPs of  
18 the same class are often capable of replacing the deleted one. The class A PBP1A and  
19 PBP1B in *E. coli* that have transglycosylase as well as transpeptidase activity can each,  
20 but not simultaneously, be deleted without any morphological consequences and are  
21 apparently not that specific [163]. Surprisingly, all of the four class A PBPs (PBP1,  
22 PBP2c, PBP2d, and PBP4) of *B. subtilis* can be simultaneously deleted resulting only in a  
23 reduction of growth rate and minor morphological effects [164]. Analysis of the

1 mucopeptide composition of the quadruple-class A mutant strain indicates that a novel  
2 unidentified enzyme must perform the glycosyltransferase activity required for  
3 peptidoglycan synthesis [164].

4 An alternative approach to elucidate the function of the various PBPs is to  
5 determine where they localize in the bacterial cell using either fluorescent protein fusions  
6 or antibodies, and fluorescence microscopy. Promising results have been obtained over  
7 the last few years (see for an comprehensive list of localized PBPs; [165]). Localization  
8 of PBPs has been studied most extensively in *B. subtilis* where 13 out of the 16 PBPs  
9 were localized using strains in which chromosomally encoded PBPs were replaced by  
10 GFP-PBP fusions [126, 166]. Three localization patterns could be observed: (i)  
11 localization in the cylindrical envelope and at the septum, (ii) septal localization only,  
12 and (iii) localization in form of a collection of foci that might be organized in arc-like or  
13 helical structures [126]. The last pattern belonged to PBPs that were either lowly  
14 expressed or presumably needed for specific growth or sporulation (PBP3, PBP4a, and  
15 PBP4\*). The class A PBP1B and the class B PBP2B localized at the septum. The class A  
16 PBP4, the class B PBP2a, and the major carboxypeptidase PBP5 [167] localized as well  
17 in the cylindrical wall as at the septum.

18 The for lateral growth responsible class B PBP2 of *E. coli* was also shown to be  
19 present in the cylindrical part as well as at the constriction using a functional GFP-PBP2  
20 [127] whereas the class B PBP3 responsible for septal peptidoglycan synthesis localizes  
21 specifically at the constriction [168]. In the case of PBP2 it was shown that its  
22 localization at the constriction was required to maintain the diameter of the new cell poles  
23 [127]. Therefore, it seems likely that the new cell poles of rod shaped bacteria are

1 synthesized with the lateral mode of PG synthesis as well as the septal mode of synthesis,  
2 whereas the cylindrical part is only synthesized in the lateral mode. This hypothesis is  
3 supported by the observation that some cocci change their shape or diameter after  
4 addition of the, for PBP2 specific, antibiotic mecillinam or deletion of genes encoding for  
5 proteins that are part of their putative elongation machinery ([6, 169]; see also this issue  
6 Y).

7         The combination of a class A and a class B PBP is in principle sufficient for bulk  
8 cell wall synthesis. Based on the localization studies in *B. subtilis* one could envision  
9 two separated PG synthesizing machineries as all ready proposed in the eighties by  
10 Higgings and Schockman [170]. One for lateral PG synthesis and one for septal PG  
11 synthesis where the latter is assisted by the lateral synthesizing machinery during  
12 synthesis of the new cell poles. However the observation that the Class A PBP1B  
13 localizes in the lateral wall as well as at the septum, where it directly associates with the  
14 class B PBP3 [101, 171], which localizes exclusively at the division site, suggests sooner  
15 that a rearrangement of the lateral PG machinery occurs to included PBPs such as PBP3  
16 that are specialized in septal PG synthesis. If this would indeed be the case, the, for  
17 PBP1B inter exchangeable, PBP1A is expected to localize as well at the constriction site.

18         The Low Molecular Weight PBPs modify the peptidoglycan by making the  
19 peptide side chains inaccessible for cross-linking or by cleaving crosslinks (see this issue  
20 X). The morphological defects of deletion of different combinations of the LMW PBPs of  
21 *E. coli* produces cells with random shapes, length and diameters [171, 172]. Therefore,  
22 the majority of these PBPs are presumably important for fine-tuning of the PG synthesis  
23 [173]. Especially the DD-caboxypeptidase PBP5 of *E. coli* seems to be important for the

1 maintenance of cell diameter, surface uniformity and overall topology of the  
2 peptidoglycan layer [149]. Therefore, it is not surprising that *E. coli* PBP5 localizes like  
3 *B. subtilis* PBP5 in the lateral wall and predominantly at the septum (unpublished data  
4 Karczmarek, den Blaauwen et al.). In other words it seems to be where the highest PG  
5 synthesizing activity occurs. This would agree with the high carboxypeptidase activity  
6 reported during division in synchronized cell cultures [174, 175]. In contrast, the  
7 carboxypeptidase PBP3 of *Streptococcus pneumoniae* localizes everywhere but at the site  
8 of septal peptidoglycan synthesis from which was concluded that by making the peptide  
9 side chains of the glycan strands unavailable for crosslinking, the synthesizing PBPs  
10 would not be able to localize and insert new cell wall material ([176] and this issue X).  
11 Interestingly, overproduction of PBP5 in *E. coli* results in spherical growth [177], which  
12 suggests that PBP2 is selective for pentapeptides as peptide cross bridge donors. Possibly,  
13 the number of PBP5 molecules of *E. coli* and *B. subtilis* determines the substrate  
14 availability of PBP2 during lateral growth and constriction/septation. Elevated levels of  
15 PBP5 or 6 can restore cell division in *fisI23*(Ts) mutants at the restrictive temperature.  
16 Larger amounts of DD-carboxypeptidase increase the amount of tetrapeptide and  
17 tripeptide acceptors preferentially used by PBP3. A similar correction of the mutation can  
18 be performed by the addition at low concentration of D-cycloserine, which inhibits the D-  
19 Ala-D-Ala ligase and thus increases the amount of tripeptide. The level of LD-  
20 carboxypeptidase that hydrolyses the tetrapeptide into tripeptide fluctuates during the cell  
21 cycle with a sharp increase at the time of division [178]. The proportion of penta-, tetra-  
22 and tripeptide might thus be an important factor in determining the cell shape by  
23 controlling the relative rate of division and elongation [99].

## **6. Attachment of the outer membrane to peptidoglycan and consequences for elongation and division**

### **6.1. Linkage between the outer membrane and the petidoglycan**

The peptidoglycan layer is connected to the outer membrane via a number of proteins. Of these, the Braun's lipoprotein (Lpp) is the only known protein that is covalently bound to the peptidoglycan while inserted in the outer membrane by its acyl chain [179]. Lpp is the most abundant outer membrane protein and exists in a free (60-70 %) and to peptidoglycan bound (30-40%) form [180, 181]. The L,D transpeptidase YbiS, and to a lesser extent the proteins ErfK and YcfS, are responsible for the linkage of the  $\alpha$ -carbonyl of A<sub>2</sub>pm to the side chain amide of the L-Lys residue located at the C-terminus of Lpp [160]. The free form might stabilize the peptidoglycan bound form by forming trimers [182]. Lpp is homogenously distributed along the cell envelope [183, 184] and by its sheer numbers, 100,000 copies per  $\mu\text{m}^2$  [185], it is in general essential for the maintenance of contacts between the outer membrane and the peptidoglycan layer.

Other protein like the peptidoglycan-associated lipoprotein (PAL) and outer membrane proteins A, C, and F (OmpA, OmpC, and OmpF) interact non-covalently with the sacculus [186, 187]. Deletion of either Lpp or Pal causes loss in the integrity of the outer membrane, release of periplasmic proteins and outer membrane vesicle formation or blebbing [188, 189]. Overexpression of Pal can compensate for the loss of Lpp in an *lpp* deletion strain, but overexpression of Lpp cannot rescue a *pal* deletion strain [190], suggesting that Pal has a more specific function than Lpp.

1 Pal is anchored to the outer membrane by a N-diacyl glyceride moiety at its  
2 second serine residue [191, 192] and interacts strongly with the peptide moiety of the  
3 peptidoglycan layer by its carboxy-terminal region [193, 194]). TolB competes with the  
4 peptidoglycan peptide side chain for the same region on Pal [195] TolB is part of the Tol-  
5 Pal system that connects the inner membrane with the outer membrane. It consists of at  
6 least the cytoplasmic membrane proteins TolA, TolQ, and TolR that interact by their  
7 transmembrane helices [196], and the periplasmic TolB and Pal protein anchored to the  
8 outer membrane. TolQ is an integral membrane protein with three membrane helices  
9 whereas TolR and TolA are bitopic membrane proteins with two domains located in the  
10 periplasm (see for a review [197]). The C-terminal domain of TolA interacts with the N  
11 terminal domain of TonB [198, 199]. The crystal structure of TolB reveals a two-domain  
12 structure [200, 201] with a mixed  $\alpha$  helix plus sheet of 5  $\beta$  strands N-terminal domain  
13 connected to a six-bladed  $\beta$ -propeller C-terminal domain. This propeller domain interacts  
14 with Pal. In addition to the interaction of Pal with TolB and peptidoglycan, it interacts  
15 independently with OmpA and with TolA [202]. The interaction of TolA with Pal is lost  
16 after dissipation of the electrochemical gradient of ions or protons [202, 203]. Based on  
17 sequence comparisons and mutagenesis studies, TolQR have been proposed to form a  
18 proton or ion channel, which converts the proton motive force (pmf) to mechanical  
19 energy to drive a conformational change in the TolA protein [204]. The channel is  
20 probably formed by 14 transmembrane helices derived from a 4:2 stoichiometry of TolQ  
21 to TolR [205]. The C-terminal periplasmic domain of TolR seems to have a role in the  
22 regulation of the opening and closing of the channel entrance [206]. Whether it regulates

1 the ion flow by forming a penetrating loop, a plug or a surface structure at the  
2 periplasmic side of the channel is not yet clear.

3 All five proteins of the Tol-Pal system accumulate at the division site during the  
4 constriction process and mutants lacking this system exhibit a delay in outer membrane  
5 invagination and contain large membrane blebs at the constriction site and the cell poles  
6 [207]. The localization of the Tol-Pal at mid cell is dependent on the presence of FtsN  
7 and therefore, probably on a mature and functioning divisome. TolA which normally  
8 requires FtsN for its recruitment to the divisome, localizes in a *ftsN* null strain suggesting  
9 that FtsN does not recruit the Tol-Pal complex directly [107]. Gerding et al., [207]  
10 propose that the trans-envelope Tol-Pal system constitutes a subcomplex of the division  
11 machinery in Gram-negative bacteria that is specifically employed to ensure proper  
12 invagination of the, to the peptidoglycan layer linked, outer membrane. The Tol-pal  
13 proteins appear to be uniformly distributed in the bacterial envelope of young cells, to  
14 accumulate at mid cell and follow the leading edge of constriction in dividing cells. This  
15 dynamic behavior is only possible if the transperiplasmic connection is of a transient  
16 nature. The competition of TonB and PG for the same binding site on Pal, and maybe of  
17 TolB and TolA for Pal, might allow a discontinuous connection that enables the Pal and  
18 Ton proteins to migrate to mid cell. As the same time the connection is sufficiently  
19 continuous to force the outer membrane to follow the invaginating cytoplasmic  
20 membrane and PG layer and even to be resistant to plasmolysis [208].

21 The pmf dependence of the Tol-pal interaction might also be related to the  
22 frequent occurrence of cell division protein mutants such as FtsZ84(TS) that are sensitive  
23 to low or high osmotic strength. Tol-Pal mutant themselves form chains of cells under

osmotic stress conditions [207, 209] indicating that they are deficient in cell division. The cell division proteins FtsE/X have been characterized as conditional osmoremedial essential proteins and deletion of their genes could be compensated for by overproduction of FtsQAZ or FtsN [79]. Reddy [79] suggested that the assembly of the divisome is possibly intrinsically osmosensitive. This osmosensitivity might be caused by the pmf dependence of the Tol-Pal system.

## **6.2 Mobility of outer membrane proteins is constrained at cell poles**

The strong and complex interaction between OM and murein sacculus has a particularly dramatic manifestation in the behaviour of OM proteins at the cell poles. Label and chase experiments designed to follow segregation and mobility of OM proteins showed that proteins at the poles were essentially immobile, whilst in other locations they became progressively intermixed with newly synthesized proteins in the course of cell elongation [210]. In vivo studies on the dynamics of the OM protein LamB also demonstrated coexistence of mobile and static populations of the protein [211]. A likely reason for such unexpected behaviour lays in the inert nature of the peptidoglycan (IPG) making up the polar caps of the sacculus [3]. Since peptidoglycan subunits in the polar regions are locked at their positions, mobility of any other (macro)molecules interacting with these affixed subunits would be constrained. In the case of covalently bound proteins as Braun's lipoprotein [160, 180], or strongly interacting with the sacculus as OmpA [212] or the Tol-Pal complex [207], the proteins become effectively anchored to the subjacent IPG. Because these proteins do interact with the OM, and are present in large numbers, they can in turn affect mobility of other OM proteins and even the physico



1 chemical properties of the polar area as such. Indeed, differentiation of ectopic or  
2 abnormal IPG regions in sacculi of certain mutants is systematically associated to a  
3 parallel development of "low protein mobility" areas in the OM [141, 150], thus  
4 reinforcing a causal relationship. Whether every single protein species in the OM is  
5 similarly affected in the polar area is unknown. Applied methods preferentially labelled a  
6 set of proteins accessible to reagents in the extracellular space [210, 211]. Therefore, free  
7 movement of non-reacting protein species in and out of the polar regions cannot be  
8 excluded. Polar IPG regions might not only influence OM protein mobility but also  
9 secretion. Fluorescently labelled proteins in the poles show a very slow decline in  
10 absolute signal intensity with time, indicating a similarly slow rate of dilution with newly  
11 made, unlabeled proteins [210]. Although still a controversial matter [213, 214], diffusion  
12 of periplasmic materials could also be influenced by the temporal stability of the OM-  
13 sacculus complex in the polar regions.

14         The constraint of OM protein mobility at the cell poles is likely to play important  
15 roles in a number of cellular functions, as the generation of asymmetries in the  
16 distribution of cell envelope proteins [215], polar targeting of specific proteins as IcsA  
17 [216, 217], secretion of autotransported proteins [218], twitching mobility [219] and  
18 possibly others.

## 20 **7. L-forms**

21  
22 The definition of what should be called an L-form has been discussed since 1939, when it  
23 was shown that many bacterial species gave rise to forms similar to the L1 culture (L

1 from Lister Institute) that Klieneberger had isolated from *Streptobacillus moniliformis*.  
2 Since 1942 the methodology for establishing L-forms routinely involves numerous  
3 passages on complex hypertonic penicillin plates over an extended period of time. The  
4 first growing cells obtained, the unstable L-forms, are spherical and osmosensitive, and  
5 they revert to normal morphology in the absence of penicillin. After further passages,  
6 often extending over several years, stable (non-reverting) derivatives are obtained, and  
7 some authors have suggested that only these should be called L-forms [220]. Others,  
8 however, have presented convincing evidence that stabilization is a secondary event,  
9 which simply prevents the reconstitution of a normal cell wall in the absence of penicillin  
10 but does not affect L-form growth [221]. What then is an L-form? In the absence of a  
11 recognized authority empowered to establish such definitions, the wisest course is to  
12 describe clearly the origin and cultivation of the organisms used, whatever name they go  
13 by. This, unfortunately, is not always the case in the L-form literature. To avoid  
14 confusion, we will call the cefsulodine or penicillin-induced spherical and osmosensitive  
15 forms, as “L-form-like” cells.

16 Because of the prolonged growth of the L-forms in the presence of penicillin and  
17 their mycoplasma-like morphology, they are thought to be devoid of cell wall. Electron  
18 microscopy showed in some cases that there is no visible cell wall in L-forms of various  
19 bacterial species, including *E. coli* [222]. Biochemical analyses of cell wall constituents  
20 in L-forms have given variable results, with numerous reports in which muramic acid,  
21 A<sub>2</sub>pm, D-glutamate, or glucosamine was or was not detected in extracts of L-forms of  
22 various bacteria, usually with little quantification [222-224]. So, we are unaware of any

1 published data that eliminate the possibility of a residual peptidoglycan in an established  
2 L-form.

3 D'Ari and coworkers [225] have found that the growth of L-form-like *E. coli* cells  
4 induced by cefsulodin or penicillin, requires residual peptidoglycan synthesis amounting,  
5 in the former case, to 7% of that of wild type cells. This is in some ways reminiscent of  
6 the paradox of chlamydial species, which are reputed to have no cell wall yet seem to  
7 require peptidoglycan synthesis, probably for cell division [226]. The amount of  
8 peptidoglycan in L-form-like cells is far too little to form a sacculus covering the entire  
9 cell [227]. Although techniques are not presently available for locating this peptidoglycan  
10 within the cell, the following arguments suggest that it may be at the division site of the  
11 spherical L-form-like cells. When, the for the septation process specific, PBP3 is  
12 inhibited by piperacillin in cefsulodin-induced L-form-like cells, a rapid block of the  
13 viable cell count is observed. This suggests that PBP3 is required for the propagation of  
14 the L-form-Like cells. The transglycosylase activity of PBP1B is required for rapid  
15 growth of the L-form-like cells, and this protein has been shown to interact directly with  
16 PBP3 [101]. In addition, PBP1B has been implicated in cell division under certain  
17 conditions [228]. The division protein FtsZ is also required for the propagation of L-  
18 form-like cells [225]. The simplest hypothesis to explain these observations –  
19 requirements for peptidoglycan synthesis, FtsZ, PBP3, and PBP1B – is that cell division  
20 in cefsulodin-induced L-form-like cells, as in rods, by means of peptidoglycan  
21 synthesized in the division plane and indispensable for cytokinesis [225].

22 It was recently reported that an established *E. coli* L-form isolated nearly 40 years  
23 ago has acquired mutations in several genes required for peptidoglycan synthesis and cell

1 division [229]. From the sequence of 36 kb of L-form DNA, the authors deduced that the  
2 FtsA, FtsW and MurG proteins have one or two amino acid changes each, the *ftsQ* gene  
3 has an amber triplet at codon 132 (of 276 codons), and the *mraY* gene has a frame shift in  
4 codon 294 that should produce a protein of 298 amino acids (instead of 360). The  
5 functional consequences, in rod-shaped cells, of the missense mutations and of the  
6 truncation of MraY are unknown. The truncated FtsQ protein would almost certainly be  
7 non-functional for cell division in rod-shaped cells, although a low level of amber  
8 suppression could provide the 22 molecules of intact FtsQ estimated to be needed for  
9 division [230]. Further characterization of this classical L-form should establish clearly  
10 whether or not the cells carry out residual peptidoglycan synthesis and, if they do,  
11 whether it is essential for their propagation.

12 Cell division in normal bacteria is inseparable from cell wall synthesis at the  
13 septum, and now [225] it has been shown that this requirement is also present for L-form-  
14 like cells. The question is why cell division would require peptidoglycan synthesis, and  
15 the simplest answer is that macromolecular PG is both substrate and product of the  
16 reaction, and the residual 7% of macromolecular PG would provide the required scaffold  
17 for assembly of division leading edge.

## 18 19 **Concluding remarks**

20 The elongation of the sacculus is performed by a dynamic protein complex “the  
21 elongase” that inserts peptidoglycan precursors at a limited number of discrete sites while  
22 using the cytoskeletal MreB helix as tracking device. The elongase consists of at least;  
23 MraY, MurG, MreBCD, RodA, PBP1, and PBP2. Based on the average number of MurG

1 foci in the cylindrical part of the cell [46] and the number of PBP2 molecules per average  
2 cell [231], it seems unrealistic to expect more than 50 elongases per average cell  
3 depending on the growth rate (Fig 9). Upon initiation of cell division by positioning of  
4 the cytoskeletal Z-ring at mid cell, a switch from the dispersed to a concentrated local  
5 peptidoglycan synthesis occurs. From this point on peptidoglycan synthesis is for a large  
6 part redirected from the elongating activity to the synthesis of new cell poles that consists  
7 of new peptidoglycan. This synthesis is performed by the divisome or septosome.  
8 Initially the synthesis seems to be performed by a premature divisome that contains PIPS  
9 (see section 4.4) to synthesize new peptidoglycan. After about one fifth of a generation  
10 time the divisome matures and might be envisioned as an extended elongase because  
11 apart from its basic peptidoglycan synthesis activity specific functions have to be added.  
12 These are, the change in morphology from cylinder to sphere (e.g. FtsK, FtsQLB, FtsW,  
13 PBP3), the invagination of the outer membrane (e.g. Tol-Pal-FtsN), and the hydrolases  
14 that allow the extension and invagination of the peptidoglycan layer (e.g. MipA, MltA,  
15 AmiA, EnvA). Again, based on the number of tightly membrane bound FtsZ molecules  
16 (unpublished data den Blaauwen) and the FtsQ, PBP3 molecules present per average cell  
17 [90, 231], not more than 50 divisome subassemblies will be present at the division site  
18 (Fig. 9). As the rate of peptidoglycan synthesis at the leading edge is constant during cell  
19 division, these subassemblies will be active during the progression of the leading edge to  
20 the closure of the constriction into a separated daughter cell pole (Fig. 9). The elongase  
21 and the divisome are dynamic hyperstructures [232] that probably at least share part of  
22 their proteins (e.g. PBP1, PBP2, MurG, MraY?, carboxypeptidases). A picture is  
23 emerging in which the PBPs, on which the focus has been for the major part of the last

century, are the proteins that are the least involved in the regulation of the creation of the shape maintaining sacculus.

## Acknowledgements

We thank Benoit Wolf for the drawing of figure 2 and 3 and Elsevier Limited Oxford, UK for the permission to reproduce figure 4 (here figure 6) from reference xx. This work was supported in part by the European commission within the 'EUR-INTAFAR' LSHM-CT-2004-512138, and 'COBRA' LSHM-CT-2003-503335 networks and by a Vernieuwingsimpuls grant 016.001.024 of the Netherlands Organization for Scientific Research (NWO) to T.d.B., by the Belgian State, Prime Minister's Office, Science Policy programming (IAP no. P6/19), the *Actions de Recherche Concertées* (grant 03/08-297), the *Fond de la Recherche Fondamentale Collective* (contrat 2.4543.05) to M. N., and BFU2006-04574 from *Ministerio de Educación y Ciencia*, Spain to J.A.A. and an institutional grant from *Fundación Ramón Areces*.

## References

- [1] Young, K.D. (2003) Bacterial shape. *Mol Microbiol* 49, 571-580.
- [2] Matsuzawa, H., Asoh, S., Kunai, K., Muraiso, K., Takasuga, A. and Ohta, T. (1989) Nucleotide sequence of the rodA gene, responsible for the rod shape of *Escherichia coli*: rodA and the pbpA gene, encoding penicillin-binding protein 2, constitute the rodA operon. *J Bacteriol* 171, 558-560.
- [3] Matsuzawa, H., Hayakawa, K., Sato, T. and Imahori, K. (1973) Characterization and genetic analysis of a mutant of *Escherichia coli* K-12 with rounded morphology. *J Bacteriol* 115, 436-442.
- [4] Henriques, A.O., Glaser, P., Piggot, P.J. and Moran, C.P., Jr. (1998) Control of cell shape and elongation by the rodA gene in *Bacillus subtilis*. *Mol Microbiol* 28, 235-247.
- [5] Costa, C.S. and Anton, D.N. (1999) Conditional lethality of cell shape mutations of *Salmonella typhimurium*: rodA and mre mutants are lethal on solid but not in liquid medium. *Curr Microbiol* 38, 137-142.
- [6] Thibessard, A., Fernandez, A., Gintz, B., Leblond-Bourget, N. and Decaris, B. (2002) Effects of rodA and pbp2b disruption on cell morphology and oxidative stress response of *Streptococcus thermophilus* CNRZ368. *J Bacteriol* 184, 2821-2826.
- [7] Ishino, F., Park, W., Tomioka, S., Tamaki, S., Takase, I., Kunugita, K., Matsuzawa, H., Asoh, S., Ohta, T., Spratt, B.G. and et al. (1986) Peptidoglycan synthetic activities in membranes of *Escherichia coli* caused by overproduction of penicillin-binding protein 2 and rodA protein. *J Biol Chem* 261, 7024-7031.

- [8] van Dam, V., Sijbrandi, R., Kol, M., Swiezewska, E., de Kruijff, B. and Breukink, E. (2007) Transmembrane transport of peptidoglycan precursors across model and bacterial membranes. *Mol Microbiol* 64, 1105-1114.
- [9] Lund, F. and Tybring, L. (1972) 6 -amidinopenicillanic acids--a new group of antibiotics. *Nat New Biol* 236, 135-137.
- [10] Spratt, B.G. and Pardee, A.B. (1975) Penicillin-binding proteins and cell shape in *E. coli*. *Nature* 254, 516-517.
- [11] Spratt, B.G. (1975) Distinct penicillin binding proteins involved in the division, elongation, and shape of *Escherichia coli* K12. *Proc Natl Acad Sci U S A* 72, 2999-3003.
- [12] Vinella, D., Joseleau-Petit, D., Thevenet, D., Boulloc, P. and D'Ari, R. (1993) Penicillin-binding protein 2 inactivation in *Escherichia coli* results in cell division inhibition, which is relieved by FtsZ overexpression. *J Bacteriol* 175, 6704-6710.
- [13] Vinella, D., D'Ari, R., Jaffe, A. and Boulloc, P. (1992) Penicillin binding protein 2 is dispensable in *Escherichia coli* when ppGpp synthesis is induced. *Embo J* 11, 1493-1501.
- [14] Joseleau-Petit, D., Thevenet, D. and D'Ari, R. (1994) ppGpp concentration, growth without PBP2 activity, and growth-rate control in *Escherichia coli*. *Mol Microbiol* 13, 911-917.
- [15] Costa, C.S. and Anton, D.N. (1993) Round-cell mutants of *Salmonella typhimurium* produced by transposition mutagenesis: lethality of *rodA* and *mre* mutations. *Mol Gen Genet* 236, 387-394.
- [16] Wachi, M. and Matsuhashi, M. (1989) Negative control of cell division by *mreB*, a gene that functions in determining the rod shape of *Escherichia coli* cells. *J Bacteriol* 171, 3123-3127.
- [17] Kruse, T., Bork-Jensen, J. and Gerdes, K. (2005) The morphogenetic MreBCD proteins of *Escherichia coli* form an essential membrane-bound complex. *Mol Microbiol* 55, 78-89.
- [18] Formstone, A. and Errington, J. (2005) A magnesium-dependent *mreB* null mutant: implications for the role of *mreB* in *Bacillus subtilis*. *Mol Microbiol* 55, 1646-1657.
- [19] Alaedini, A. and Day, R.A. (1999) Identification of two penicillin-binding multienzyme complexes in *Haemophilus influenzae*. *Biochem Biophys Res Commun* 264, 191-195.
- [20] Normark, S. (1969) Mutation in *Escherichia coli* K-12 mediating spherulike envelopes and changes tolerance to ultraviolet irradiation and some antibiotics. *J Bacteriol* 98, 1274-1277.
- [21] Wachi, M., Doi, M., Okada, Y. and Matsuhashi, M. (1989) New *mre* genes *mreC* and *mreD*, responsible for formation of the rod shape of *Escherichia coli* cells. *J Bacteriol* 171, 6511-6516.
- [22] Wachi, M., Doi, M., Tamaki, S., Park, W., Nakajima-Iijima, S. and Matsuhashi, M. (1987) Mutant isolation and molecular cloning of *mre* genes, which determine cell shape, sensitivity to mecillinam, and amount of penicillin-binding proteins in *Escherichia coli*. *J Bacteriol* 169, 4935-4940.
- [23] Jones, L.J., Carballido-Lopez, R. and Errington, J. (2001) Control of cell shape in bacteria: helical, actin-like filaments in *Bacillus subtilis*. *Cell* 104, 913-922.
- [24] Shih, Y.L., Le, T. and Rothfield, L. (2003) Division site selection in *Escherichia coli* involves dynamic redistribution of Min proteins within coiled structures that extend between the two cell poles. *Proc Natl Acad Sci U S A* 100, 7865-7870.

- [25] Figge, R.M., Divakaruni, A.V. and Gober, J.W. (2004) MreB, the cell shape-determining bacterial actin homologue, co-ordinates cell wall morphogenesis in *Caulobacter crescentus*. *Mol Microbiol* 51, 1321-1332.
- [26] van den Ent, F., Amos, L.A. and Lowe, J. (2001) Prokaryotic origin of the actin cytoskeleton. *Nature* 413, 39-44.
- [27] Esue, O., Cordero, M., Wirtz, D. and Tseng, Y. (2005) The assembly of MreB, a prokaryotic homolog of actin. *J Biol Chem* 280, 2628-2635.
- [28] Karimova, G., Pidoux, J., Ullmann, A. and Ladant, D. (1998) A bacterial two-hybrid system based on a reconstituted signal transduction pathway. *Proc Natl Acad Sci U S A* 95, 5752-5756.
- [29] van den Ent, F., Leaver, M., Bendezu, F., Errington, J., de Boer, P. and Lowe, J. (2006) Dimeric structure of the cell shape protein MreC and its functional implications. *Mol Microbiol* 62, 1631-1642.
- [30] Karczmarek, A., Martínez-Arteaga Baselga, R., Alexeeva, S., Hansen, F.G., Vicente, M., Nanninga, N. and den Blaauwen, T. (2007) DNA and origin region segregation are not affected by the transition from rod to sphere after inhibition of *Escherichia coli* MreB by A22. *Mol. Microbiol.* 65, 51-63.
- [31] Iwai, N., Nagai, K. and Wachi, M. (2002) Novel S-benzylisothiourea compound that induces spherical cells in *Escherichia coli* probably by acting on a rod-shape-determining protein(s) other than penicillin-binding protein 2. *Biosci Biotechnol Biochem* 66, 2658-2662.
- [32] Iwai, N., Ebata, T., Nagura, H., Kitazume, T., Nagai, K. and Wachi, M. (2004) Structure-activity relationship of S-benzylisothiourea derivatives to induce spherical cells in *Escherichia coli*. *Biosci Biotechnol Biochem* 68, 2265-2269.
- [33] Varma, A., de Pedro, M. and Young, K.D. (2007) FtsZ directs a second mode of peptidoglycan synthesis in *Escherichia coli*. *J Bacteriol.*
- [34] Divakaruni, A.V., Loo, R.R., Xie, Y., Loo, J.A. and Gober, J.W. (2005) The cell-shape protein MreC interacts with extracytoplasmic proteins including cell wall assembly complexes in *Caulobacter crescentus*. *Proc Natl Acad Sci U S A* 102, 18602-18607.
- [35] Campo, N., Tjalsma, H., Buist, G., Stepniak, D., Meijer, M., Veenhuis, M., Westermann, M., Muller, J.P., Bron, S., Kok, J., Kuipers, O.P. and Jongbloed, J.D. (2004) Subcellular sites for bacterial protein export. *Mol Microbiol* 53, 1583-1599.
- [36] Dye, N.A., Pincus, Z., Theriot, J.A., Shapiro, L. and Gitai, Z. (2005) Two independent spiral structures control cell shape in *Caulobacter*. *Proc Natl Acad Sci U S A* 102, 18608-18613.
- [37] Leaver, M. and Errington, J. (2005) Roles for MreC and MreD proteins in helical growth of the cylindrical cell wall in *Bacillus subtilis*. *Mol Microbiol* 57, 1196-1209.
- [38] Walker, S., Chen, L., Hu, Y., Rew, Y., Shin, D. and Boger, D.L. (2005) Chemistry and biology of ramoplanin: a lipoglycopeptide with potent antibiotic activity. *Chem Rev* 105, 449-476.
- [39] Fang, X., Tiyanont, K., Zhang, Y., Wanner, J., Boger, D. and Walker, S. (2006) The mechanism of action of ramoplanin and enduracidin. *Mol Biosyst* 2, 69-76.
- [40] Tiyanont, K., Doan, T., Lazarus, M.B., Fang, X., Rudner, D.Z. and Walker, S. (2006) Imaging peptidoglycan biosynthesis in *Bacillus subtilis* with fluorescent antibiotics. *Proc Natl Acad Sci U S A* 103, 11033-11038.



- [41] Kim, S.Y., Gitai, Z., Kinkhabwala, A., Shapiro, L. and Moerner, W.E. (2006) Single molecules of the bacterial actin MreB undergo directed treadmilling motion in *Caulobacter crescentus*. *Proc Natl Acad Sci U S A* 103, 10929-10934.
- [42] Defeu Soufo, H.J. and Graumann, P.L. (2006) Dynamic localization and interaction with other *Bacillus subtilis* actin-like proteins are important for the function of MreB. *Mol Microbiol* 62, 1340-1356.
- [43] Mengin-Lecreulx, D., Texier, L., Rousseau, M. and van Heijenoort, J. (1991) The murG gene of *Escherichia coli* codes for the UDP-N-acetylglucosamine: N-acetylmuramyl-(pentapeptide) pyrophosphoryl-undecaprenol N-acetylglucosamine transferase involved in the membrane steps of peptidoglycan synthesis. *J Bacteriol* 173, 4625-4636.
- [44] Chen, L., Men, H., Ha, S., Ye, X.Y., Brunner, L., Hu, Y. and Walker, S. (2002) Intrinsic lipid preferences and kinetic mechanism of *Escherichia coli* MurG. *Biochemistry* 41, 6824-6833.
- [45] Ha, S., Walker, D., Shi, Y. and Walker, S. (2000) The 1.9 Å crystal structure of *Escherichia coli* MurG, a membrane-associated glycosyltransferase involved in peptidoglycan biosynthesis. *Protein Sci* 9, 1045-1052.
- [46] Mohammadi, T., Karczmarek, A., Crouvoisier, M., Bouhss, A., Mengin-Lecreulx, D. and den Blaauwen, T. (2007) The essential peptidoglycan glycosyltransferase MurG forms a complex with proteins involved in lateral envelope growth as well as the proteins involved in cell division in *Escherichia coli*. *Mol. Microbiol.* in press.
- [47] Ikeda, M., Wachi, M., Jung, H.K., Ishino, F. and Matsubashi, M. (1991) The *Escherichia coli* mraY gene encoding UDP-N-acetylmuramoyl-pentapeptide: undecaprenyl-phosphate phospho-N-acetylmuramoyl-pentapeptide transferase. *J Bacteriol* 173, 1021-1026.
- [48] Boyle, D.S. and Donachie, W.D. (1998) mraY is an essential gene for cell growth in *Escherichia coli*. *J Bacteriol* 180, 6429-6432.
- [49] Bouhss, A., Mengin-Lecreulx, D., Le Beller, D. and Van Heijenoort, J. (1999) Topological analysis of the MraY protein catalysing the first membrane step of peptidoglycan synthesis. *Mol Microbiol* 34, 576-585.
- [50] Brandish, P.E., Burnham, M.K., Lonsdale, J.T., Southgate, R., Inukai, M. and Bugg, T.D. (1996) Slow binding inhibition of phospho-N-acetylmuramyl-pentapeptide-translocase (*Escherichia coli*) by mureidomycin A. *J Biol Chem* 271, 7609-7614.
- [51] Stachyra, T., Dini, C., Ferrari, P., Bouhss, A., van Heijenoort, J., Mengin-Lecreulx, D., Blanot, D., Biton, J. and Le Beller, D. (2004) Fluorescence detection-based functional assay for high-throughput screening for MraY. *Antimicrob Agents Chemother* 48, 897-902.
- [52] Aaron, M., Charbon, G., Lam, H., Schwarz, H., Vollmer, W. and Jacobs-Wagner, C. (2007) The tubulin homologue FtsZ contributes to cell elongation by guiding cell wall precursor synthesis in *Caulobacter crescentus*. *Mol Microbiol* 64, 938-952.
- [53] Oliva, M.A., Cordell, S.C. and Lowe, J. (2004) Structural insights into FtsZ protofilament formation. *Nat Struct Mol Biol* 11, 1243-1250.
- [54] Pla, J., Sanchez, M., Palacios, P., Vicente, M. and Aldea, M. (1991) Preferential cytoplasmic location of FtsZ, a protein essential for *Escherichia coli* septation. *Mol Microbiol* 5, 1681-1686.

- [55] Anderson, D.E., Gueiros-Filho, F.J. and Erickson, H.P. (2004) Assembly dynamics of FtsZ rings in *Bacillus subtilis* and *Escherichia coli* and effects of FtsZ-regulating proteins. *J Bacteriol* 186, 5775-5781.
- [56] Chen, Y. and Erickson, H.P. (2005) Rapid in vitro assembly dynamics and subunit turnover of FtsZ demonstrated by fluorescence resonance energy transfer. *J Biol Chem* 280, 22549-22554.
- [57] Harry, E., Monahan, L. and Thompson, L. (2006) Bacterial cell division: the mechanism and its precision. *Int Rev Cytol* 253, 27-94.
- [58] Thanedar, S. and Margolin, W. (2004) FtsZ exhibits rapid movement and oscillation waves in helix-like patterns in *Escherichia coli*. *Curr Biol* 14, 1167-1173.
- [59] Peters, P.C., Migocki, M.D., Thoni, C. and Harry, E.J. (2007) A new assembly pathway for the cytokinetic Z ring from a dynamic helical structure in vegetatively growing cells of *Bacillus subtilis*. *Mol Microbiol* 64, 487-499.
- [60] Addinall, S.G. and Lutkenhaus, J. (1996) FtsA is localized to the septum in an FtsZ-dependent manner. *J Bacteriol* 178, 7167-7172.
- [61] Hale, C.A. and de Boer, P.A. (1999) Recruitment of ZipA to the septal ring of *Escherichia coli* is dependent on FtsZ and independent of FtsA. *J Bacteriol* 181, 167-176.
- [62] Pichoff, S. and Lutkenhaus, J. (2007) Identification of a region of FtsA required for interaction with FtsZ. *Mol Microbiol* 64, 1129-1138.
- [63] Hale, C.A. and de Boer, P.A. (1997) Direct binding of FtsZ to ZipA, an essential component of the septal ring structure that mediates cell division in *E. coli*. *Cell* 88, 175-185.
- [64] van den Ent, F. and Lowe, J. (2000) Crystal structure of the cell division protein FtsA from *Thermotoga maritima*. *Embo J* 19, 5300-5307.
- [65] Pichoff, S. and Lutkenhaus, J. (2005) Tethering the Z ring to the membrane through a conserved membrane targeting sequence in FtsA. *Mol Microbiol* 55, 1722-1734.
- [66] Mileykovskaya, E. and Dowhan, W. (2000) Visualization of phospholipid domains in *Escherichia coli* by using the cardiolipin-specific fluorescent dye 10-N-nonyl acridine orange. *J Bacteriol* 182, 1172-1175.
- [67] Koppelman, C.M., Den Blaauwen, T., Duursma, M.C., Heeren, R.M. and Nanninga, N. (2001) *Escherichia coli* minicell membranes are enriched in cardiolipin. *J Bacteriol* 183, 6144-6147.
- [68] Mileykovskaya, E. and Dowhan, W. (2005) Role of membrane lipids in bacterial division-site selection. *Curr Opin Microbiol* 8, 135-142.
- [69] van den Brink-van der Laan, E., Boots, J.W., Spelbrink, R.E., Kool, G.M., Breukink, E., Killian, J.A. and de Kruijff, B. (2003) Membrane interaction of the glycosyltransferase MurG: a special role for cardiolipin. *J Bacteriol* 185, 3773-3779.
- [70] Geissler, B. and Margolin, W. (2005) Evidence for functional overlap among multiple bacterial cell division proteins: compensating for the loss of FtsK. *Mol Microbiol* 58, 596-612.
- [71] Geissler, B., Shiomi, D. and Margolin, W. (2007) The *ftsA\** gain-of-function allele of *Escherichia coli* and its effects on the stability and dynamics of the Z ring. *Microbiology* 153, 814-825.
- [72] Rico, A.I., Garcia-Ovalle, M., Mingorance, J. and Vicente, M. (2004) Role of two essential domains of *Escherichia coli* FtsA in localization and progression of the division ring. *Mol Microbiol* 53, 1359-1371.

- [73] Low, H.H., Moncrieffe, M.C. and Lowe, J. (2004) The crystal structure of ZapA and its modulation of FtsZ polymerisation. *J Mol Biol* 341, 839-852.
- [74] Small, E., Marrington, R., Rodger, A., Scott, D.J., Sloan, K., Roper, D., Dafforn, T.R. and Addinall, S.G. (2007) FtsZ polymer-bundling by the *Escherichia coli* ZapA orthologue, YgfE, involves a conformational change in bound GTP. *J Mol Biol* 369, 210-221.
- [75] Levin, P.A., Kurtser, I.G. and Grossman, A.D. (1999) Identification and characterization of a negative regulator of FtsZ ring formation in *Bacillus subtilis*. *Proc Natl Acad Sci U S A* 96, 9642-9647.
- [76] Chung, K.M., Hsu, H.H., Yeh, H.Y. and Chang, B.Y. (2007) Mechanism of regulation of prokaryotic tubulin-like GTPase FtsZ by membrane protein EzrA. *J Biol Chem* 282, 14891-14897.
- [77] Schmidt, K.L., Peterson, N.D., Kustusch, R.J., Wissel, M.C., Graham, B., Phillips, G.J. and Weiss, D.S. (2004) A predicted ABC transporter, FtsEX, is needed for cell division in *Escherichia coli*. *J Bacteriol* 186, 785-793.
- [78] Corbin, B.D., Wang, Y., Beuria, T.K. and Margolin, W. (2007) Interaction between cell division proteins FtsE and FtsZ. *J Bacteriol* 189, 3026-3035.
- [79] Reddy, M. (2007) Role of FtsEX in cell division of *Escherichia coli*: viability of ftsEX mutants is dependent on functional SufI or high osmotic strength. *J Bacteriol* 189, 98-108.
- [80] Aarsman, M.E., Piette, A., Fraipont, C., Vinkenvleugel, T.M., Nguyen-Disteche, M. and den Blaauwen, T. (2005) Maturation of the *Escherichia coli* divisome occurs in two steps. *Mol Microbiol* 55, 1631-1645.
- [81] Wang, X., Possoz, C. and Sherratt, D.J. (2005) Dancing around the divisome: asymmetric chromosome segregation in *Escherichia coli*. *Genes Dev* 19, 2367-2377.
- [82] Wang, S., Arends, S.J., Weiss, D.S. and Newman, E.B. (2005) A deficiency in S-adenosylmethionine synthetase interrupts assembly of the septal ring in *Escherichia coli* K-12. *Mol Microbiol* 58, 791-799.
- [83] Hara, H., Yasuda, S., Horiuchi, K. and Park, J.T. (1997) A promoter for the first nine genes of the *Escherichia coli* mra cluster of cell division and cell envelope biosynthesis genes, including ftsI and ftsW. *J Bacteriol* 179, 5802-5811.
- [84] Bigot, S., Sivanathan, V., Possoz, C., Barre, F.X. and Cornet, F. (2007) FtsK, a literate chromosome segregation machine. *Mol Microbiol* 64, 1434-1441.
- [85] Dorazi, R. and Dewar, S.J. (2000) Membrane topology of the N-terminus of the *Escherichia coli* FtsK division protein. *FEBS Lett* 478, 13-18.
- [86] Begg, K.J., Dewar, S.J. and Donachie, W.D. (1995) A new *Escherichia coli* cell division gene, ftsK. *J Bacteriol* 177, 6211-6222.
- [87] Robson, S.A. and King, G.F. (2006) Domain architecture and structure of the bacterial cell division protein DivIB. *Proc Natl Acad Sci U S A* 103, 6700-6705.
- [88] Sanchez-Pulido, L., Devos, D., Genevrois, S., Vicente, M. and Valencia, A. (2003) POTRA: a conserved domain in the FtsQ family and a class of beta-barrel outer membrane proteins. *Trends Biochem Sci* 28, 523-526.
- [89] Scheffers, D.-J., Robichon, C., Haan, G.J., den Blaauwen, T., Koningstein, G., van Bloois, E. and Luirink, J. (2007) Role of the FtsQ transmembrane segment in localization to the cell division site. *J. Bacteriol.* submitted.

- [90] D'Ulisse, V., Fagioli, M., Ghelardini, P. and Paolozzi, L. (2007) Three functional subdomains of the *Escherichia coli* FtsQ protein are involved in its interaction with the other division proteins. *Microbiology* 153, 124-138.
- [91] Di Lallo, G., Fagioli, M., Barionovi, D., Ghelardini, P. and Paolozzi, L. (2003) Use of a two-hybrid assay to study the assembly of a complex multicomponent protein machinery: bacterial septosome differentiation. *Microbiology* 149, 3353-3359.
- [92] Karimova, G., Dautin, N. and Ladant, D. (2005) Interaction network among *Escherichia coli* membrane proteins involved in cell division as revealed by bacterial two-hybrid analysis. *J Bacteriol* 187, 2233-2243.
- [93] Dai, K. and Lutkenhaus, J. (1992) The proper ratio of FtsZ to FtsA is required for cell division to occur in *Escherichia coli*. *J Bacteriol* 174, 6145-6151.
- [94] Piette, A., Fraipont, C., Den Blaauwen, T., Aarsman, M.E., Pastoret, S. and Nguyen-Disteche, M. (2004) Structural determinants required to target penicillin-binding protein 3 to the septum of *Escherichia coli*. *J Bacteriol* 186, 6110-6117.
- [95] Buddelmeijer, N., Judson, N., Boyd, D., Mekalanos, J.J. and Beckwith, J. (2002) YgbQ, a cell division protein in *Escherichia coli* and *Vibrio cholerae*, localizes in codependent fashion with FtsL to the division site. *Proc Natl Acad Sci U S A* 99, 6316-6321.
- [96] Goehring, N.W., Gueiros-Filho, F. and Beckwith, J. (2005) Premature targeting of a cell division protein to midcell allows dissection of divisome assembly in *Escherichia coli*. *Genes Dev* 19, 127-137.
- [97] Buddelmeijer, N. and Beckwith, J. (2004) A complex of the *Escherichia coli* cell division proteins FtsL, FtsB and FtsQ forms independently of its localization to the septal region. *Mol Microbiol* 52, 1315-1327.
- [98] Goehring, N.W., Gonzalez, M.D. and Beckwith, J. (2006) Premature targeting of cell division proteins to midcell reveals hierarchies of protein interactions involved in divisome assembly. *Mol Microbiol* 61, 33-45.
- [99] Begg, K.J., Takasuga, A., Edwards, D.H., Dewar, S.J., Spratt, B.G., Adachi, H., Ohta, T., Matsuzawa, H. and Donachie, W.D. (1990) The balance between different peptidoglycan precursors determines whether *Escherichia coli* cells will elongate or divide. *J Bacteriol* 172, 6697-6703.
- [100] Adam, M., Fraipont, C., Rhazi, N., Nguyen-Disteche, M., Lakaye, B., Frere, J.M., Devreese, B., Van Beeumen, J., van Heijenoort, Y., van Heijenoort, J. and Ghuysen, J.M. (1997) The bimodular G57-V577 polypeptide chain of the class B penicillin-binding protein 3 of *Escherichia coli* catalyzes peptide bond formation from thioesters and does not catalyze glycan chain polymerization from the lipid II intermediate. *J Bacteriol* 179, 6005-6009.
- [101] Bertsche, U., Kast, T., Wolf, B., Fraipont, C., Aarsman, M.E., Kannenberg, K., von Rechenberg, M., Nguyen-Disteche, M., den Blaauwen, T., Holtje, J.V. and Vollmer, W. (2006) Interaction between two murein (peptidoglycan) synthases, PBP3 and PBP1B, in *Escherichia coli*. *Mol Microbiol* 61, 675-690.
- [102] Schiffer, G. and Holtje, J.V. (1999) Cloning and characterization of PBP 1C, a third member of the multimodular class A penicillin-binding proteins of *Escherichia coli*. *J Biol Chem* 274, 32031-32039.
- [103] Vollmer, W., von Rechenberg, M. and Holtje, J.V. (1999) Demonstration of molecular interactions between the murein polymerase PBP1B, the lytic transglycosylase

MltA, and the scaffolding protein MipA of *Escherichia coli*. *J Biol Chem* 274, 6726-6734.

[104] Yang, J.C., Van Den Ent, F., Neuhaus, D., Brevier, J. and Lowe, J. (2004) Solution structure and domain architecture of the divisome protein FtsN. *Mol Microbiol* 52, 651-660.

[105] Goehring, N.W., Robichon, C. and Beckwith, J. (2007) Role for the nonessential N terminus of FtsN in divisome assembly. *J Bacteriol* 189, 646-649.

[106] Corbin, B.D., Geissler, B., Sadasivam, M. and Margolin, W. (2004) Z-ring-independent interaction between a subdomain of FtsA and late septation proteins as revealed by a polar recruitment assay. *J Bacteriol* 186, 7736-7744.

[107] Bernard, C.S., Sadasivam, M., Shiomi, D. and Margolin, W. (2007) An altered FtsA can compensate for the loss of essential cell division protein FtsN in *Escherichia coli*. *Mol Microbiol* 64, 1289-1305.

[108] Bernhardt, T.G. and de Boer, P.A. (2003) The *Escherichia coli* amidase AmiC is a periplasmic septal ring component exported via the twin-arginine transport pathway. *Mol Microbiol* 48, 1171-1182.

[109] Heidrich, C., Templin, M.F., Ursinus, A., Merdanovic, M., Berger, J., Schwarz, H., de Pedro, M.A. and Holtje, J.V. (2001) Involvement of N-acetylmuramyl-L-alanine amidases in cell separation and antibiotic-induced autolysis of *Escherichia coli*. *Mol Microbiol* 41, 167-178.

[110] Bernhardt, T.G. and de Boer, P.A. (2004) Screening for synthetic lethal mutants in *Escherichia coli* and identification of EnvC (YibP) as a periplasmic septal ring factor with murein hydrolase activity. *Mol Microbiol* 52, 1255-1269.

[111] Varma, A. and Young, K.D. (2004) FtsZ collaborates with penicillin binding proteins to generate bacterial cell shape in *Escherichia coli*. *J Bacteriol* 186, 6768-6774.

[112] Satta, G., Canepari, P., Maurici, R. and Marcialis, M.A. (1985) Interactions between lateral wall elongation and septum formation during cell cycle in *Klebsiella pneumoniae*. *Ann Inst Pasteur Microbiol* 136A, 85-89.

[113] Holtje, J.V. (1998) Growth of the stress-bearing and shape-maintaining murein sacculus of *Escherichia coli*. *Microbiol Mol Biol Rev* 62, 181-203.

[114] Nanninga, N. (1998) Morphogenesis of *Escherichia coli*. *Microbiol Mol Biol Rev* 62, 110-129.

[115] Errington, J., Daniel, R.A. and Scheffers, D.J. (2003) Cytokinesis in bacteria. *Microbiol Mol Biol Rev* 67, 52-65, table of contents.

[116] Van Tubergen, R.P. and Setlow, R.B. (1961) Quantitative radioautographic studies on exponentially growing cultures of *Escherichia coli*. The distribution of parental DNA, RNA, protein, and cell wall among progeny cells. *Biophys J* 1, 589-625.

[117] Ryter, A., Hirota, Y. and Schwarz, U. (1973) Process of cellular division in *Escherichia coli* growth pattern of *E. coli* murein. *J Mol Biol* 78, 185-195.

[118] de Chastellier, C., Hellio, R. and Ryter, A. (1975) Study of cell wall growth in *Bacillus megaterium* by high-resolution autoradiography. *J Bacteriol* 123, 1184-1196.

[119] Woldringh, C.L., Huls, P., Pas, E., Brakenhoff, G.J. and Nanninga, N. (1987) Topography of peptidoglycan synthesis during elongation and polar cap formation in a cell division mutant of *Escherichia coli* MC4100. *J. Gen. Microbiol.* 133, 575-586.

- [120] Mulder, E. and Woldringh, C.L. (1991) Autoradiographic analysis of diaminopimelic acid incorporation in filamentous cells of *Escherichia coli*: repression of peptidoglycan synthesis around the nucleoid. *J Bacteriol* 173, 4751-4756.
- [121] Wientjes, F.B. and Nanninga, N. (1989) Rate and topography of peptidoglycan synthesis during cell division in *Escherichia coli*: concept of a leading edge. *J Bacteriol* 171, 3412-3419.
- [122] Burman, L.G., Raichler, J. and Park, J.T. (1983) Evidence for diffuse growth of the cylindrical portion of the *Escherichia coli* murein sacculus. *J Bacteriol* 155, 983-988.
- [123] Burman, L.G., Reichler, J. and Park, J.T. (1983) Evidence for multisite growth of *Escherichia coli* murein involving concomitant endopeptidase and transpeptidase activities. *J Bacteriol* 156, 386-392.
- [124] Cooper, S., Hsieh, M.L. and Guenther, B. (1988) Mode of peptidoglycan synthesis in *Salmonella typhimurium*: single-strand insertion. *J Bacteriol* 170, 3509-3512.
- [125] de Pedro, M.A., Holtje, J.V. and Schwarz, H. (2002) Fast lysis of *Escherichia coli* filament cells requires differentiation of potential division sites. *Microbiology* 148, 79-86.
- [126] Scheffers, D.J., Jones, L.J. and Errington, J. (2004) Several distinct localization patterns for penicillin-binding proteins in *Bacillus subtilis*. *Mol Microbiol* 51, 749-764.
- [127] Den Blaauwen, T., Aarsman, M.E., Vischer, N.O. and Nanninga, N. (2003) Penicillin-binding protein PBP2 of *Escherichia coli* localizes preferentially in the lateral wall and at mid-cell in comparison with the old cell pole. *Mol Microbiol* 47, 539-547.
- [128] Daniel, R.A. and Errington, J. (2003) Control of cell morphogenesis in bacteria: two distinct ways to make a rod-shaped cell. *Cell* 113, 767-776.
- [129] de Pedro, M.A., Quintela, J.C., Holtje, J.V. and Schwarz, H. (1997) Murein segregation in *Escherichia coli*. *J Bacteriol* 179, 2823-2834.
- [130] Burman, L.G. and Park, J.T. (1984) Molecular model for elongation of the murein sacculus of *Escherichia coli*. *Proc Natl Acad Sci U S A* 81, 1844-1848.
- [131] Spratt, B.G. (1977) Properties of the penicillin-binding proteins of *Escherichia coli* K12. *Eur J Biochem* 72, 341-352.
- [132] Cooper, S. and Hsieh, M.L. (1988) The rate and topography of cell wall synthesis during the division cycle of *Escherichia coli* using N-acetylglucosamine as a peptidoglycan label. *J Gen Microbiol* 134, 1717-1721.
- [133] Born, P., Breukink, E. and Vollmer, W. (2006) In vitro synthesis of cross-linked murein and its attachment to sacculi by PBP1A from *Escherichia coli*. *J Biol Chem* 281, 26985-26993.
- [134] Bertsche, U., Breukink, E., Kast, T. and Vollmer, W. (2005) In vitro murein peptidoglycan synthesis by dimers of the bifunctional transglycosylase-transpeptidase PBP1B from *Escherichia coli*. *J Biol Chem* 280, 38096-38101.
- [135] Shih, Y.L. and Rothfield, L. (2006) The bacterial cytoskeleton. *Microbiol Mol Biol Rev* 70, 729-754.
- [136] Koch, A.L. and De Pedro, M.A. (2006) Partition of old murein in small patches over the entire wall of *E. coli* cells forced to grow as a coccoid. *Curr Microbiol* 52, 249-253.
- [137] De Pedro, M.A., Schwarz, H. and Koch, A.L. (2003) Patchiness of murein insertion into the sidewall of *Escherichia coli*. *Microbiology* 149, 1753-1761.

- [138] Vollmer, W. and Holtje, J.V. (2001) Morphogenesis of *Escherichia coli*. *Curr Opin Microbiol* 4, 625-633.
- [139] Schwarz, U., Asmus, A. and Frank, H. (1969) Autolytic enzymes and cell division of *Escherichia coli*. *J Mol Biol* 41, 419-429.
- [140] Nanninga, N. (1991) Cell division and peptidoglycan assembly in *Escherichia coli*. *Mol Microbiol* 5, 791-795.
- [141] de Pedro, M.A., Donachie, W.D., Holtje, J.V. and Schwarz, H. (2001) Constitutive septal murein synthesis in *Escherichia coli* with impaired activity of the morphogenetic proteins RodA and penicillin-binding protein 2. *J Bacteriol* 183, 4115-4126.
- [142] Caparros, M., Pisabarro, A.G. and de Pedro, M.A. (1992) Effect of D-amino acids on structure and synthesis of peptidoglycan in *Escherichia coli*. *J Bacteriol* 174, 5549-5559.
- [143] Romeis, T., Kamper, J. and Kahmann, R. (1997) Single-chain fusions of two unrelated homeodomain proteins trigger pathogenicity in *Ustilago maydis*. *Proc Natl Acad Sci U S A* 94, 1230-1234.
- [144] Ishidate, K., Ursinus, A., Holtje, J.V. and Rothfield, L. (1998) Analysis of the length distribution of murein glycan strands in *ftsZ* and *ftsI* mutants of *E. coli*. *FEMS Microbiol Lett* 168, 71-75.
- [145] Holtje, J.V. and Heidrich, C. (2001) Enzymology of elongation and constriction of the murein sacculus of *Escherichia coli*. *Biochimie* 83, 103-108.
- [146] Glauner, B. and Holtje, J.V. (1990) Growth pattern of the murein sacculus of *Escherichia coli*. *J Biol Chem* 265, 18988-18996.
- [147] Pisabarro, A.G., de Pedro, M.A. and Vazquez, D. (1985) Structural modifications in the peptidoglycan of *Escherichia coli* associated with changes in the state of growth of the culture. *J Bacteriol* 161, 238-242.
- [148] Nelson, D.E. and Young, K.D. (2001) Contributions of PBP 5 and DD-carboxypeptidase penicillin binding proteins to maintenance of cell shape in *Escherichia coli*. *J Bacteriol* 183, 3055-3064.
- [149] Nelson, D.E. and Young, K.D. (2000) Penicillin binding protein 5 affects cell diameter, contour, and morphology of *Escherichia coli*. *J Bacteriol* 182, 1714-1721.
- [150] de Pedro, M.A., Young, K.D., Holtje, J.V. and Schwarz, H. (2003) Branching of *Escherichia coli* cells arises from multiple sites of inert peptidoglycan. *J Bacteriol* 185, 1147-1152.
- [151] Nanninga, N., Wientjes, F.B., de Jonge, B.L. and Woldringh, C.L. (1990) Polar cap formation during cell division in *Escherichia coli*. *Res Microbiol* 141, 103-118.
- [152] Heidrich, C., Ursinus, A., Berger, J., Schwarz, H. and Holtje, J.V. (2002) Effects of multiple deletions of murein hydrolases on viability, septum cleavage, and sensitivity to large toxic molecules in *Escherichia coli*. *J Bacteriol* 184, 6093-6099.
- [153] Priyadarshini, R., de Pedro, M.A. and Young, K.D. (2007) Role of peptidoglycan amidases in the development and morphology of the division septum in *Escherichia coli*. *J Bacteriol*.
- [154] Priyadarshini, R., Popham, D.L. and Young, K.D. (2006) Daughter cell separation by penicillin-binding proteins and peptidoglycan amidases in *Escherichia coli*. *J Bacteriol* 188, 5345-5355.

- [155] Olijhoek, A.J., Klencke, S., Pas, E., Nanninga, N. and Schwarz, U. (1982) Volume growth, murein synthesis, and murein cross-linkage during the division cycle of *Escherichia coli* PA3092. *J Bacteriol* 152, 1248-1254.
- [156] Den Blaauwen, T., Buddelmeijer, N., Aarsman, M.E., Hameete, C.M. and Nanninga, N. (1999) Timing of FtsZ assembly in *Escherichia coli*. *J Bacteriol* 181, 5167-5175.
- [157] Wientjes, F.B. and Nanninga, N. (1991) On the role of the high molecular weight penicillin-binding proteins in the cell cycle of *Escherichia coli*. *Res Microbiol* 142, 333-344.
- [158] von Rechenberg, M., Ursinus, A. and Holtje, J.V. (1996) Affinity chromatography as a means to study multienzyme complexes involved in murein synthesis. *Microb Drug Resist* 2, 155-157.
- [159] Di Berardino, M., Dijkstra, A., Stuber, D., Keck, W. and Gubler, M. (1996) The monofunctional glycosyltransferase of *Escherichia coli* is a member of a new class of peptidoglycan-synthesising enzymes. *FEBS Lett* 392, 184-188.
- [160] Magnet, S., Bellais, S., Dubost, L., Fourgeaud, M., Mainardi, J.L., Petit-Frere, S., Marie, A., Mengin-Lecreulx, D., Arthur, M. and Gutmann, L. (2007) Identification of the L,D-Transpeptidases Responsible for Attachment of the Braun Lipoprotein to *Escherichia coli* Peptidoglycan. *J Bacteriol* 189, 3927-3931.
- [161] Botta, G.A. and Park, J.T. (1981) Evidence for involvement of penicillin-binding protein 3 in murein synthesis during septation but not during cell elongation. *J Bacteriol* 145, 333-340.
- [162] Popham, D.L. and Young, K.D. (2003) Role of penicillin-binding proteins in bacterial cell morphogenesis. *Curr Opin Microbiol* 6, 594-599.
- [163] Yousif, S.Y., Broome-Smith, J.K. and Spratt, B.G. (1985) Lysis of *Escherichia coli* by beta-lactam antibiotics: deletion analysis of the role of penicillin-binding proteins 1A and 1B. *J Gen Microbiol* 131, 2839-2845.
- [164] McPherson, D.C. and Popham, D.L. (2003) Peptidoglycan synthesis in the absence of class A penicillin-binding proteins in *Bacillus subtilis*. *J Bacteriol* 185, 1423-1431.
- [165] Scheffers, D.J. and Pinho, M.G. (2005) Bacterial cell wall synthesis: new insights from localization studies. *Microbiol Mol Biol Rev* 69, 585-607.
- [166] Scheffers, D.J. and Errington, J. (2004) PBP1 is a component of the *Bacillus subtilis* cell division machinery. *J Bacteriol* 186, 5153-5156.
- [167] Lawrence, P.J. and Strominger, J.L. (1970) Biosynthesis of the peptidoglycan of bacterial cell walls. XVI. The reversible fixation of radioactive penicillin G to the D-alanine carboxypeptidase of *Bacillus subtilis*. *J Biol Chem* 245, 3660-3666.
- [168] Weiss, D.S., Chen, J.C., Ghigo, J.M., Boyd, D. and Beckwith, J. (1999) Localization of FtsI (PBP3) to the septal ring requires its membrane anchor, the Z ring, FtsA, FtsQ, and FtsL. *J Bacteriol* 181, 508-520.
- [169] Pinho, M.G., de Lencastre, H. and Tomasz, A. (2000) Cloning, characterization, and inactivation of the gene *pbpC*, encoding penicillin-binding protein 3 of *Staphylococcus aureus*. *J Bacteriol* 182, 1074-1079.
- [170] Higgins, M.L. and Shockman, G.D. (1971) Procaryotic cell division with respect to wall and membranes. *CRC Crit Rev Microbiol* 1, 29-72.
- [171] Denome, S.A., Elf, P.K., Henderson, T.A., Nelson, D.E. and Young, K.D. (1999) *Escherichia coli* mutants lacking all possible combinations of eight penicillin binding



proteins: viability, characteristics, and implications for peptidoglycan synthesis. J  
 Bacteriol 181, 3981-3993.  
 [172] Vega, D. and Ayala, J.A. (2006) The DD-carboxypeptidase activity encoded by  
 pbp4B is not essential for the cell growth of *Escherichia coli*. Arch Microbiol 185, 23-27.  
 [173] Nilsen, T., Ghosh, A.S., Goldberg, M.B. and Young, K.D. (2004) Branching sites  
 and morphological abnormalities behave as ectopic poles in shape-defective *Escherichia*  
*coli*. Mol Microbiol 52, 1045-1054.  
 [174] Mirelman, D., Yashouv-Gan, Y., Nuchamovitz, Y., Rozenhak, S. and Ron, E.Z.  
 (1978) Murein biosynthesis during a synchronous cell cycle of *Escherichia coli* B. J  
 Bacteriol 134, 458-461.  
 [175] Mirelman, D., Yashouv-Gan, Y. and Schwarz, U. (1977) Regulation of murein  
 biosynthesis and septum formation in filamentous cells of *Escherichia coli* PAT 84. J  
 Bacteriol 129, 1593-1600.  
 [176] Morlot, C., Noirclerc-Savoye, M., Zapun, A., Dideberg, O. and Vernet, T. (2004)  
 The D,D-carboxypeptidase PBP3 organizes the division process of *Streptococcus*  
*pneumoniae*. Mol Microbiol 51, 1641-1648.  
 [177] Markiewicz, Z., Broome-Smith, J.K., Schwarz, U. and Spratt, B.G. (1982)  
 Spherical *E. coli* due to elevated levels of D-alanine carboxypeptidase. Nature 297, 702-  
 704.  
 [178] Beck, B.D. and Park, J.T. (1976) Activity of three murein hydrolases during the  
 cell division cycle of *Escherichia coli* K-12 as measured in toluene-treated cells. J  
 Bacteriol 126, 1250-1260.  
 [179] Braun, V. and Rehn, K. (1969) Chemical characterization, spatial distribution and  
 function of a lipoprotein (murein-lipoprotein) of the *E. coli* cell wall. The specific effect  
 of trypsin on the membrane structure. Eur J Biochem 10, 426-438.  
 [180] Braun, V. (1975) Covalent lipoprotein from the outer membrane of *Escherichia*  
*coli*. Biochim Biophys Acta 415, 335-377.  
 [181] Inouye, M., Shaw, J. and Shen, C. (1972) The assembly of a structural lipoprotein  
 in the envelope of *Escherichia coli*. J Biol Chem 247, 8154-8159.  
 [182] Choi, D.S., Yamada, H., Mizuno, T. and Mizushima, S. (1986) Trimeric structure  
 and localization of the major lipoprotein in the cell surface of *Escherichia coli*. J Biol  
 Chem 261, 8953-8957.  
 [183] Hiemstra, H., de Hoop, M.J., Inouye, M. and Witholt, B. (1986) Induction kinetics  
 and cell surface distribution of *Escherichia coli* lipoprotein under lac promoter control. J  
 Bacteriol 168, 140-151.  
 [184] Hiemstra, H., Nanninga, N., Woldringh, C.L., Inouye, M. and Witholt, B. (1987)  
 Distribution of newly synthesized lipoprotein over the outer membrane and the  
 peptidoglycan sacculus of an *Escherichia coli* lac-lpp strain. J Bacteriol 169, 5434-5444.  
 [185] Nikaido, H. (1996) Outer membrane, 29-47 pp. American Society for  
 Microbiology, Washington D.C.  
 [186] Koebnik, R. (1995) Proposal for a peptidoglycan-associating alpha-helical motif in  
 the C-terminal regions of some bacterial cell-surface proteins. Mol Microbiol 16, 1269-  
 1270.  
 [187] Leduc, M., Ishidate, K., Shakibai, N. and Rothfield, L. (1992) Interactions of  
*Escherichia coli* membrane lipoproteins with the murein sacculus. J Bacteriol 174, 7982-  
 7988.

[188] Suzuki, H., Nishimura, Y., Yasuda, S., Nishimura, A., Yamada, M. and Hirota, Y. (1978) Murein-lipoprotein of *Escherichia coli*: a protein involved in the stabilization of bacterial cell envelope. *Mol Gen Genet* 167, 1-9.

[189] Yem, D.W. and Wu, H.C. (1978) Physiological characterization of an *Escherichia coli* mutant altered in the structure of murein lipoprotein. *J Bacteriol* 133, 1419-1426.

[190] Cascales, E., Bernadac, A., Gavioli, M., Lazzaroni, J.C. and Lloubes, R. (2002) Pal lipoprotein of *Escherichia coli* plays a major role in outer membrane integrity. *J Bacteriol* 184, 754-759.

[191] Mizuno, T. (1979) A novel peptidoglycan-associated lipoprotein found in the cell envelope of *Pseudomonas aeruginosa* and *Escherichia coli*. *J Biochem (Tokyo)* 86, 991-1000.

[192] Gennity, J.M. and Inouye, M. (1991) The protein sequence responsible for lipoprotein membrane localization in *Escherichia coli* exhibits remarkable specificity. *J Biol Chem* 266, 16458-16464.

[193] Lazzaroni, J.C. and Portalier, R. (1992) The *excC* gene of *Escherichia coli* K-12 required for cell envelope integrity encodes the peptidoglycan-associated lipoprotein (PAL). *Mol Microbiol* 6, 735-742.

[194] Parsons, L.M., Lin, F. and Orban, J. (2006) Peptidoglycan recognition by Pal, an outer membrane lipoprotein. *Biochemistry* 45, 2122-2128.

[195] Ray, M.C., Germon, P., Vianney, A., Portalier, R. and Lazzaroni, J.C. (2000) Identification by genetic suppression of *Escherichia coli* TolB residues important for TolB-Pal interaction. *J Bacteriol* 182, 821-824.

[196] Journet, L., Rigal, A., Lazdunski, C. and Benedetti, H. (1999) Role of TolR N-terminal, central, and C-terminal domains in dimerization and interaction with TolA and tolQ. *J Bacteriol* 181, 4476-4484.

[197] Lazzaroni, J.C., Dubuisson, J.F. and Vianney, A. (2002) The Tol proteins of *Escherichia coli* and their involvement in the translocation of group A colicins. *Biochimie* 84, 391-397.

[198] Dubuisson, J.F., Vianney, A. and Lazzaroni, J.C. (2002) Mutational analysis of the TolA C-terminal domain of *Escherichia coli* and genetic evidence for an interaction between TolA and TolB. *J Bacteriol* 184, 4620-4625.

[199] Walburger, A., Lazdunski, C. and Corda, Y. (2002) The Tol/Pal system function requires an interaction between the C-terminal domain of TolA and the N-terminal domain of TolB. *Mol Microbiol* 44, 695-708.

[200] Abergel, C., Bouveret, E., Claverie, J.M., Brown, K., Rigal, A., Lazdunski, C. and Benedetti, H. (1999) Structure of the *Escherichia coli* TolB protein determined by MAD methods at 1.95 Å resolution. *Structure* 7, 1291-1300.

[201] Carr, S., Penfold, C.N., Bamford, V., James, R. and Hemmings, A.M. (2000) The structure of TolB, an essential component of the tol-dependent translocation system, and its protein-protein interaction with the translocation domain of colicin E9. *Structure* 8, 57-66.

[202] Cascales, E. and Lloubes, R. (2004) Deletion analyses of the peptidoglycan-associated lipoprotein Pal reveals three independent binding sequences including a TolA box. *Mol Microbiol* 51, 873-885.

- [203] Cascales, E., Gavioli, M., Sturgis, J.N. and Lloubes, R. (2000) Proton motive force drives the interaction of the inner membrane TolA and outer membrane pal proteins in *Escherichia coli*. *Mol Microbiol* 38, 904-915.
- [204] Goemaere, E.L., Cascales, E. and Lloubes, R. (2007) Mutational analyses define helix organization and key residues of a bacterial membrane energy-transducing complex. *J Mol Biol* 366, 1424-1436.
- [205] Cascales, E., Lloubes, R. and Sturgis, J.N. (2001) The TolQ-TolR proteins energize TolA and share homologies with the flagellar motor proteins MotA-MotB. *Mol Microbiol* 42, 795-807.
- [206] Goemaere, E.L., Devert, A., Lloubes, R. and Cascales, E. (2007) Movements of the TolR C-terminal domain depend on TolQR ionizable key residues and regulate activity of the Tol complex. *J Biol Chem*.
- [207] Gerding, M.A., Ogata, Y., Pecora, N.D., Niki, H. and de Boer, P.A. (2007) The trans-envelope Tol-Pal complex is part of the cell division machinery and required for proper outer-membrane invagination during cell constriction in *E. coli*. *Mol Microbiol* 63, 1008-1025.
- [208] MacAllister, T.J., Cook, W.R., Weigand, R. and Rothfield, L.I. (1987) Membrane-murein attachment at the leading edge of the division septum: a second membrane-murein structure associated with morphogenesis of the gram-negative bacterial division septum. *J Bacteriol* 169, 3945-3951.
- [209] Dubuisson, J.F., Vianney, A., Hugouvieux-Cotte-Pattat, N. and Lazzaroni, J.C. (2005) Tol-Pal proteins are critical cell envelope components of *Erwinia chrysanthemi* affecting cell morphology and virulence. *Microbiology* 151, 3337-3347.
- [210] de Pedro, M.A., Grunfelder, C.G. and Schwarz, H. (2004) Restricted Mobility of Cell Surface Proteins in the Polar Regions of *Escherichia coli*. *J Bacteriol* 186, 2594-2602.
- [211] Gibbs, K.A., Isaac, D.D., Xu, J., Hendrix, R.W., Silhavy, T.J. and Theriot, J.A. (2004) Complex spatial distribution and dynamics of an abundant *Escherichia coli* outer membrane protein, LamB. *Mol Microbiol* 53, 1771-1783.
- [212] Wang, Y. (2002) The function of OmpA in *Escherichia coli*. *Biochem Biophys Res Commun* 292, 396-401.
- [213] Foley, M., Brass, J.M., Birmingham, J., Cook, W.R., Garland, P.B., Higgins, C.F. and Rothfield, L.I. (1989) Compartmentalization of the periplasm at cell division sites in *Escherichia coli* as shown by fluorescence photobleaching experiments. *Mol Microbiol* 3, 1329-1336.
- [214] Mullineaux, C.W., Nenninger, A., Ray, N. and Robinson, C. (2006) Diffusion of green fluorescent protein in three cell environments in *Escherichia coli*. *J Bacteriol* 188, 3442-3448.
- [215] Lybarger, S.R. and Maddock, J.R. (2001) Polarity in action: asymmetric protein localization in bacteria. *J Bacteriol* 183, 3261-3267.
- [216] Janakiraman, A. and Goldberg, M.B. (2004) Evidence for polar positional information independent of cell division and nucleoid occlusion. *Proc Natl Acad Sci U S A* 101, 835-840.
- [217] Charles, M., Perez, M., Kobil, J.H. and Goldberg, M.B. (2001) Polar targeting of *Shigella* virulence factor IcsA in *Enterobacteriaceae* and *Vibrio*. *Proc Natl Acad Sci U S A* 98, 9871-9876.

- [218] Jain, S., van Ulsen, P., Benz, I., Schmidt, M.A., Fernandez, R., Tommassen, J. and Goldberg, M.B. (2006) Polar localization of the autotransporter family of large bacterial virulence proteins. *J Bacteriol* 188, 4841-4850.
- [219] Skerker, J.M. and Berg, H.C. (2001) Direct observation of extension and retraction of type IV pili. *Proc Natl Acad Sci U S A* 98, 6901-6904.
- [220] Klieneberger-Nobel, E. (1960) L-forms of bacteria, pp. Academic Press, New York.
- [221] Lederberg, J. and St Clair, J. (1958) Protoplasts and L-type growth of *Escherichia coli*. *J Bacteriol* 75, 143-160.
- [222] Gumpert, J. and Taubeneck, U. (1983) Characteristic properties and biological significance of stable protoplast type L-forms. *Experientia Suppl* 46, 227-241.
- [223] Gumpert, J. and Taubeneck, U. (1976) Characterization of a stable spheroplast type L-form of *Proteus mirabilis* D 52 as cell envelope mutant. II. Electronmicroscopic investigations. *Z Allg Mikrobiol* 16, 9-26.
- [224] Gumpert, J. and Taubeneck, U. (1975) Characterization of a stable spheroplast type L-form of *Proteus mirabilis* D 52 as cell envelope mutant. I. Isolation, growth characteristics, biochemical activities, and sensitivity to bacteriophages. *Z Allg Mikrobiol* 15, 399-410.
- [225] Joseleau-Petit, D., Liébart, J.-C., Ayala, J.A. and D'Ari, R. (2007) Unstable *Escherichia coli* L-forms revisited: growth requires peptidoglycan synthesis. *J. bacteriol* submitted.
- [226] McCoy, A.J. and Maurelli, A.T. (2006) Building the invisible wall: updating the chlamydial peptidoglycan anomaly. *Trends Microbiol* 14, 70-77.
- [227] Wientjes, F.B., Woldringh, C.L. and Nanninga, N. (1991) Amount of peptidoglycan in cell walls of gram-negative bacteria. *J Bacteriol* 173, 7684-7691.
- [228] Garcia del Portillo, F., de Pedro, M.A., Joseleau-Petit, D. and D'Ari, R. (1989) Lytic response of *Escherichia coli* cells to inhibitors of penicillin-binding proteins 1a and 1b as a timed event related to cell division. *J Bacteriol* 171, 4217-4221.
- [229] Siddiqui, R.A., Hoischen, C., Holst, O., Heinze, I., Schlott, B., Gumpert, J., Diekmann, S., Grosse, F. and Platzer, M. (2006) The analysis of cell division and cell wall synthesis genes reveals mutationally inactivated *ftsQ* and *mraY* in a protoplast-type L-form of *Escherichia coli*. *FEMS Microbiol Lett* 258, 305-311.
- [230] Carson, M.J., Barondess, J. and Beckwith, J. (1991) The *FtsQ* protein of *Escherichia coli*: membrane topology, abundance, and cell division phenotypes due to overproduction and insertion mutations. *J Bacteriol* 173, 2187-2195.
- [231] Dougherty, T.J., Kennedy, K., Kessler, R.E. and Pucci, M.J. (1996) Direct quantitation of the number of individual penicillin-binding proteins per cell in *Escherichia coli*. *J Bacteriol* 178, 6110-6115.
- [232] Norris, V., den Blaauwen, T., Cabin-Flaman, A., Doi, R.H., Harshey, R., Janniere, L., Jimenez-Sanchez, A., Jin, D.J., Levin, P.A., Mileykovskaya, E., Minsky, A., Saier, M., Jr. and Skarstad, K. (2007) Functional taxonomy of bacterial hyperstructures. *Microbiol Mol Biol Rev* 71, 230-253.